

16/09/99
JCSIS U.S. PTO

Please type a plus sign (+) inside this box ☐

PTO/SB/05 (12/97)

UTILITY PATENT APPLICATION TRANSMITTAL

(Only for new nonprovisional applications under 37 CFR 1.53(b))

Attorney Docket No. 20221Y Total Pages 94

First Named Inventor or Application Identifier

Rosen, et al.

Express Mail Label No. EI242726602US

JCSIS U.S. PTO
04/22/99
296662

APPLICATION ELEMENTS

See MPEP chapter 600 concerning utility patent application contents.

1. ☒ Fee Transmittal Form
(Submit an original, and a duplicate for fee processing)
2. ☒ Specification [Total Pages 69]
3. ☒ Drawing(s) (35 USC 113) [Total Sheets 17]
4. Oath or Declaration [Total Pages 4]
 - a. ☒ Newly executed (original or copy)
 - b. ☐ Copy from a prior application (37 CFR 1.63(d))
(for continuation/divisional with Box 14 completed)
[Note Box 5 below]
 - i. ☐ **DELETION OF INVENTOR(S)**
Signed statement attached deleting inventor(s) named in the prior application, see 37 CFR 1.63(d)(2) and 1.33 (b).
5. ☒ Incorporation By Reference (useable if Box 4b is checked). The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied under Box 4b, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.
6. Nucleotide and/or Amino Acid Sequence Submission (if applicable, all necessary)
 - a. ☐ Computer Readable Copy
 - b. ☐ Paper Copy (identical to computer copy)
 - c. ☐ Statement verifying identity of above copies

ADDRESS TO: Assistant Commissioner for
Box Patent Application
Washington, DC 20231

ACCOMPANYING APPLICATION PARTS

7. ☐ Assignment papers (cover sheet & document(s))
8. ☐ 37 CFR 3.73(b) Statement (when there is an assignee) ☐ Power of Attorney
9. ☐ Information Disclosure Statement (IDS)/PTO-1449 ☐ Copies of IDS Citations
10. ☐ Preliminary Amendment
11. ☒ Return Receipt Postcard (MPEP 503) (Should be specifically itemized)
12. ☐ Certified Copy of Priority Document(s) (if foreign priority is claimed)
13. ☐ Other:

14. If a CONTINUING APPLICATION, check appropriate box and supply the requisite information:

☐ Continuation ☐ Divisional ☒ Continuation-in-part (CIP) of prior application No. 60 / 082,643

15. CORRESPONDENCE ADDRESS

☒ Customer Number or Bar Code Label

Customer No. 000210

(Insert Customer No. or Attach bar code here)

NAME	Michael D. Yablonsky				
ADDRESS	Merck & Co., Inc., P. O. Box 2000				
CITY	Rahway	STATE	NJ	ZIP CODE	07065-0907
COUNTRY	USA	TELEPHONE	732-594-4678	FAX	732-594-4720

(computer generated form No. U6 - Merck & Co., Inc., 12/97)

EXPRESS MAIL CERTIFICATE

DATE OF DEPOSIT April 22, 1999

EXPRESS MAIL NO. EI242726602US

I HEREBY CERTIFY THAT THIS CORRESPONDENCE IS BEING DEPOSITED WITH THE UNITED STATES POSTAL SERVICE AS EXPRESS MAIL "POST OFFICE TO ADDRESSEE" ON THE ABOVE DATE IN AN ENVELOPE ADDRESSED TO ASSISTANT COMMISSIONER FOR PATENTS, BOX PATENT APPLICATION, WASHINGTON, D.C. 20231.

MAILED BY

DATE April 22, 1999

TITLE OF THE INVENTION

AUTOANTIGENIC FRAGMENTS, METHODS AND ASSAYS

CROSS-REFERENCE TO RELATED APPLICATIONS

5 Not applicable.

STATEMENT REGARDING FEDERALLY-SPONSORED R&D

This invention was made in part under Federally Sponsored Research. The U.S. Government may have certain rights in this
10 invention.

REFERENCE TO MICROFICHE APPENDIX

Not applicable.

15 FIELD OF THE INVENTION

This invention relates to the production of autoantigenic fragments from autoantigens and the uses of the autoantigenic fragments.

20 BACKGROUND OF THE INVENTION

The mature immune system of animals differentiates between self-molecules and non-self-molecules and mounts an immune response only against the latter. The immune system learns which molecules are self through constant exposure to those molecules that are
25 normally a part of the animal. Thus, the mature immune system is tolerized to the presence of molecules that are self. However, the immune system is not tolerized to molecules that are newly presented in the animal. These molecules can be antigens and thereby stimulate an immune response against them. Commonly, newly presented antigens
30 are from an extracorporeal source, such as an infection. In this case, the immune response helps to destroy the source of the antigens and thereby clear the infection from the body.

Newly presented antigens are produced *in vivo* through the degradation of cellular components. When the immune system
35 recognizes these degradation products of self molecules as "non-self"

antigens, an immune response can be mounted against them and an autoimmune disease can develop. Thus, these antigens are members of the class of molecules generally referred to as autoantigens and the antibodies produced against them are referred to as autoantibodies. For clarity herein, autoantigen is used to refer to the complete self molecule as found in the body. Autoantigenic fragment is used to refer to the degradation product of the autoantigen. Thus it is when an epitope is presented to the immune system as autoantigenic fragment that an immune response is elicited. Once elicited, the immune response can target the autoantigenic fragment, the autoantigen, or both.

Autoimmune diseases are diseases in which a specific immune response to self-molecules occurs, often leading to tissue and organ damage and dysfunction. The diseases can be organ-specific (e.g. Type I diabetes mellitus, thyroiditis, myasthenia gravis, primary biliary cirrhosis) or systemic in nature (e.g. systemic lupus erythematosus, rheumatoid arthritis, polymyositis, dermatomyositis, Sjogren's syndrome, scleroderma, and graft-vs.-host disease).

One source of autoantigenic fragments is cleavage of an autoantigen during apoptosis. Apoptosis is a morphologically and biochemically distinct form of cell death that occurs in many different cell types during a wide range of physiologic and pathologic circumstances (reviewed in (Jacobson *et al.*, 1997; Thompson, 1995; White, 1996)). Studies report that specific proteolysis catalyzed by a novel family of cysteine proteases is of critical importance in mediating apoptosis (Chinnaiyan and Dixit, 1996a; Martin and Green, 1995; Thornberry and Molineaux, 1995). These proteases (termed caspases), cleave downstream substrates after a consensus tetrapeptide sequence ending with aspartic acid (P_1). The caspases are synthesized as inactive precursors that require specific proteolytic cleavage after an aspartic acid residue for activation (reviewed in (Nicholson and Thornberry, 1997)).

Granzyme B, a serine protease found in the cytoplasmic granules of cytotoxic T lymphocytes (CTL) and natural killer (NK) cells, has a similar requirement to caspases, for aspartic acid in the substrate P_1 position (Odake *et al.*, 1991; Poe *et al.*, 1991). Studies have reported

that granzyme B plays an important role in inducing apoptotic nuclear changes in target cells during granule exocytosis induced cytotoxicity (Darmon *et al.*, 1996; Heusel *et al.*, 1994; Sarin *et al.*, 1997; Shresta *et al.*, 1995; Talanian *et al.*, 1997).

5 Granzyme B is described as catalyzing the cleavage and activation of several caspases (Chinnaiyan *et al.*, 1996b; Darmon *et al.*, 1995; Duan *et al.*, 1996; Fernandes-Alnemri *et al.*, 1996; Gu *et al.*, 1996; Martin *et al.*, 1996; Muzio *et al.*, 1996; Quan *et al.*, 1996; Sarin *et al.*, 1997; Song *et al.*, 1996a; Srinivasula *et al.*, 1996; Talanian *et al.*, 1997; Wang *et al.*, 1996). Granzyme B also initiates caspase-independent pathways which contribute to target cell death. However, while several candidates for these additional pathways exist, they remain largely undefined (Sarin *et al.*, 1997; Talanian *et al.*, 1997).

15 One candidate pathway is the direct proteolysis of death substrates by granzyme B, although efficient non-caspase cellular substrates for this protease have not yet been identified. Initial studies have indicated that the cleavage of PARP, U1-70kDa and lamin B observed during granzyme B-induced cell death is catalyzed by caspases, rather than directly by granzyme B (Darmon *et al.*, 1995; Martin *et al.*, 20 1996; Talanian *et al.*, 1997), but the effects of granzyme B on other caspase substrates *in vitro* and during granule-induced cytotoxicity have not been extensively studied.

BRIEF DESCRIPTION OF THE DRAWINGS

25 FIG. 1. Granzyme B cleaves purified DNA-PK_{CS}, NuMA, PARP and caspases 3 and 7 *in vitro* with different efficiencies. Reactions containing 0-12.5nM granzyme B (for DNA-PK_{CS} and NuMA) or 0-50 nM granzyme B (for caspases 3 and 7 and PARP) were performed as described in Example 1. Cleavage fragments were detected by 30 fluorography or immunoblotting (monoclonal antibody 18-2 was used to detect DNA-PK_{CS}). On the right side of each panel, the SDS-PAGE migration positions of the intact molecules are denoted by arrows, and the fragment sizes are indicated.

35 FIG. 2. Cleavage of autoantigens *in vitro* with purified granzyme B or caspase-3 yields different fragments. Reactions

containing 30nM purified DNA-PK_{CS} (lanes 1-3), endogenous DNA-PK_{CS} and NuMA in HeLa lysate (lanes 4-9) or [³⁵S]methionine-labeled PARP (lanes 10-12) were incubated with the following amounts of granzyme B: 1.25nM (lane 2); 12.5nM (lanes 5 & 8) or 50nM (lane 11). Similar experiments were performed with these substrates using the following amounts of caspase-3: 42pM (lanes 3 & 12) or 100pM (lanes 6 & 9). Reaction mixtures were incubated at 37°C for 15 min. (granzyme B reactions) or 60 min. (caspase-3 reactions). Note that 100nM Ac-DEVD-CHO was added when granzyme B was used to cleave DNA-PK_{CS} and NuMA in HeLa cell lysates, to prevent activation of HeLa cell caspases. Intact and cleaved NuMA and DNA-PK_{CS} were visualized by immunoblotting, and PARP was visualized by autoradiography. In the case of DNA-PK_{CS}, blots obtained using monoclonal antibodies 25-4 (directed against the C-terminus) and 18-2 (directed against the N-terminus) are shown in lanes (1-3) and (4-6), respectively.

FIG. 3. Kinase activity of DNA-PK_{CS} is abolished after cleavage with granzyme B. Kinase assays were performed using intact DNA-PK_{CS} (lanes 1 & 2) or granzyme B-cleaved DNA-PK_{CS} (lanes 3 & 4) in the absence (lanes 2 & 4) or presence (lanes 1 & 3) of 10μg/ml of DNA. DNA-PK_{CS} itself was omitted from the otherwise complete kinase reaction mix as a control in lane 5. Phosphorylation of SP1 substrate was detected by autoradiography.

FIG. 4. Endogenous DNA-PK_{CS} and NuMA in HeLa cell lysates are cleaved in a caspase-independent manner after adding purified granzyme B and incubating *in vitro*. 12.5 nM purified granzyme B (lanes 5-7) or 105 pM purified caspase-3 (lanes 2-4) were added to lysates of control HeLa cells, in the presence of 100 nM Ac-DEVD-CHO (lanes 3 & 6) or 100nM Ac-YVAD-CHO (lanes 4 & 7). After incubating at 37°C for 60 min, the reactions were terminated. DNA-PK_{CS}, NuMA and PARP were detected by immunoblotting as described in Example 1 (monoclonal antibody 18-2 was used to detect DNA-PK_{CS}). Equal amounts of protein were electrophoresed in each lane.

FIG. 5. Granzyme B cleaves DNA-PK_{CS} at VGPD2698-F2699 and DEVD2712-N2713. A [³⁵S]methionine-labeled wild-type (wt) DNA-

PK_{CS} polypeptide (DNA-PK_{CS}²⁵⁶⁶⁻²⁹²⁸) and two radiolabeled polypeptides containing mutations in the P1 positions of the predicted granzyme B cleavage site (D2698A) and the known caspase-3 cleavage site (D2712A), were generated as described in Example 1. These polypeptides were incubated in the absence of added proteases (lanes 1, 4 & 7), or in the presence of 8nM recombinant caspase-3 (lanes 2, 5 & 8) or 8 nM purified granzyme B (lanes 3, 6 & 9) for 60 min at 37°C. After terminating the reactions, samples were electrophoresed, and the intact polypeptide and the cleavage products were detected by fluorography. Fragment sizes of 18kDa (a), 28kDa (b), 20kDa (c) and 26kDa (d) were generated (see lower panel for schematic representation).

FIG. 6. Endogenous DNA-PK_{CS}, NuMA and PARP are cleaved after *in vivo* incubation of intact K562 cells with YT cell granule contents. K562 cells were incubated for 30 min at 37°C in the absence (lanes 1-3) or presence (lane 4) of 100 µM Ac-DEVD-CHO. Aliquots of these cell suspensions (each containing 3 x 10⁵ K562 cells) were then further incubated for 90 min at 37°C in the presence of 1mM Ca²⁺ (lane 1), 1 mM EDTA + YT cell granule contents (lane 2) or 1 mM Ca²⁺ + YT cell granule contents (lanes 3 & 4). After terminating the reactions, DNA-PK_{CS}, NuMA and PARP were detected by immunoblotting as described in the Example 1 (Patient serum G.A. was used to blot DNA-PK_{CS}).

FIG. 7. Granzyme B-specific fragment of DNA-PK_{CS} is generated in K562 cells attacked by LAK cells. Fas-negative K562 target cells were preincubated in the absence (lanes 1-3) or presence (lane 4) of 100µM Ac-DEVD-CHO for 1 hr, followed by co-incubation for a further 4 hr at 37°C (effector:target ratio 5:1). After terminating the reactions, the following numbers of cells were electrophoresed in each gel lane: 1.7 x 10⁶ LAK cells (lane 1); 0.34 x 10⁶ K562 cells (lane 2); 1.7 x 10⁶ LAK cells plus 0.34 x 10⁶ K562 cells (lanes 3 & 4). DNA-PK_{CS} and PARP were detected by immunoblotting; patient serum G.A. was used to detect DNA-PK_{CS}.

FIG. 8. Ac-DEVD-CHO-insensitive nuclear morphologic changes are induced in intact HeLa cells after *in vivo* incubation with

YT cell granule contents. HeLa cells were incubated with YT cell granule contents for 1hr at 37°C in the absence (8A & 8B) or presence (8C) of 100µM Ac-DEVD-CHO as described in Experimental Procedures. After fixation and permeabilization, cells were stained with antibodies to

5 PARP, as well as propidium iodide (PI) and DAPI. Antibody staining was visualized with FITC-goat anti-human antibodies. Merged images of antibody staining (green), PI staining (red) and DAPI staining (blue) are presented. (8A, 8B): YT cell granule contents induce prominent PI-rich surface blebs, nuclear condensation and fragmentation, with a

10 characteristic redistribution of PARP to the rim of the condensing nucleus (8A) or apoptotic bodies (8B, arrows). (8C): Granule contents induce marked nuclear condensation even in the presence of Ac-DEVD-CHO (arrows); an adjacent, normal HeLa cell nucleus that has not yet undergone morphologic change is shown for comparison (arrowhead).

15 Note that treatment with Ac-DEVD-CHO abolishes the formation of PI-rich surface blebs, nuclear fragmentation, and the redistribution of PARP. Size bar: 8A: 4.4 µm; 8B: 4.0 µm; 8C: 6.6 µm.

FIGS. 9A-9B shows a 2101 amino acid sequence of NuMA as found on Entrez at ACCESSION 284337; PID g284337; DBSOURCE PIR: locus A42184.

20

FIGS. 10A-10B shows a 2115 amino acid sequence of NuMA as found on Entrez at ACCESSION 107227, PID g107227; DBSOURCE PIR: locus S23647.

FIGS. 11A-11C shows the amino acid sequence of DNA PK_{CS} as found on Entrez at ACCESSION 1362789; PID g1362789; DBSOURCE PIR: locus A57099.

25

FIGS. 12A-12B shows the amino acid sequence of PARP as found on Entrez at ACCESSION 130781; PID g130781; DBSOURCE SWISS-PROT: locus PPOL_HUMAN, accession P09874 (listing only

30 references 1 & 2 of 12).

SUMMARY OF THE INVENTION

The present invention provides autoantigenic fragments and methods for their use in the treatment of autoimmune disease. Also

provided are assays for detecting an autoimmune condition in an animal, including the presence of an autoimmune disease.

Caspase-mediated proteolysis of downstream substrates is a critical element of the central execution pathway common to all forms of apoptosis studied to date. While cytolytic lymphocyte granule-induced cell death activates this caspase-dependent pathway, recent studies have also provided evidence for caspase-independent pathways in this form of cell death. However, non-caspase substrates for granzyme B (and potentially other granule proteases) during granule-induced cell death have not previously been defined. The present invention makes use of the observation that cellular components are directly and efficiently cleaved by granule contents, including in particular granzyme B, *in vitro* and *in vivo*, and that this cleavage leads to the generation of unique autoantigenic fragments not observed during other forms of apoptosis. This direct, caspase-independent ability of granzyme B to cleave downstream death substrates to autoantigenic fragments is an apoptotic effector mechanism which is insensitive to inhibitors of the signaling or execution components of the endogenous apoptotic cascade.

An aspect of this invention is a composition that includes at least one autoantigenic fragment. The autoantigenic fragment is produced by the action of a granule enzyme on an autoantigen. In a preferred embodiment, the enzyme is a granule enzyme of CTL, NK or LAK cell granules. In a most preferred embodiment, the enzyme is granzyme B and the antigenic fragment is produced by the cleavage of the autoantigen by granzyme B at a site that is not cleaved by a caspase. In a preferred embodiment the autoantigen is DNA PK_{CS}, PARP or NuMA. In a most preferred embodiment, the autoantigenic fragment is one or more of DNA-PK_{CS} from amino acid 2699 to 4096; DNA-PK_{CS} from amino acid 3211 to 4096; PARP from amino acid 1 to 537; PARP from amino acid 538 to 1004; NuMA from amino acid 412 to 2111 and NuMA from amino acid 1 to 1799.

An aspect of this invention is a pharmaceutical composition made with one or more purified and isolated autoantigenic fragments. In a preferred embodiment, the autoantigenic fragment has at least one end derived from granzyme B cleavage at a site in the autoantigen that is

not cleaved by a caspase. A pharmaceutically acceptable carrier is also included. In a preferred embodiment the composition includes one of more of the following autoantigenic fragments: DNA-PK_{CS} from amino acids 2699 to 4096; DNA-PK_{CS} from amino acids 3211 to 4096; PARP from amino acid 1 to 537; PARP from amino acids 538 to 1004; NuMA from amino acids 412 to 2111 and NuMA from amino acids 1 to 1799. In another preferred embodiment the pharmaceutical composition includes one or more autoantigenic fragments derived from a malignant cell.

An aspect of this invention is a method of treating a patient in need of treatment for an autoimmune disease by administering to the patient an autoantigenic fragment of this invention. The autoimmune disease can be organ specific, *e.g.*, Type I diabetes mellitus, thyroiditis, myasthenia gravis, primary biliary cirrhosis, or systemic in nature *e.g.* systemic lupus erythematosus, rheumatoid arthritis, polymyositis, dermatomyositis, Sjogren's syndrome, scleroderma, and graft-vs.-host disease. In one preferred embodiment, the treatment is therapeutic. For example, a patient suffering from an immune disease can be administered an autoantigenic fragment by contacting the sera of the patient with the fragment under conditions that allow the binding of autoantibodies in the sera to bind to the fragment. In this embodiment, the level of autoantibodies circulating in the patient can be reduced. In another embodiment the treatment is prophylactic. In this embodiment, a patient who is at risk of developing an autoimmune disease is tolerized to at least one autoantigenic fragment. Thereafter, the risk of, or severity of an autoimmune disease arising upon the later production of the autoantigenic fragment *in vivo*, is reduced or eliminated. In a preferred embodiment, a patient is tolerized by identifying a target tissue to which an autoimmune disease can arise, providing at least one granule enzyme, contacting the granule enzyme with cells from the target tissue to produce autoantigenic fragments of autoantigens present in the cells. The autoantigenic fragments are then administered to the patient to tolerize the patient to the presence of the fragments. In preferred embodiments, the autoantigens can be partly or wholly purified from the cells of the target tissue. The granule enzyme can also

be partly or wholly purified before contacting with the autoantigens. The enzyme can also be made by recombinant methods. In a preferred embodiment, the autoantigenic fragments are partly or wholly purified before they are administered to the patient. In prophylactic methods of tolerizing, the autoantigenic fragments are administered in pharmaceutically acceptable compositions that are designed not to raise an immune response to the fragments, *i.e.*, no immunostimulatory adjuvants are administered with the fragments. In a preferred embodiment, the treatment uses one or more of the following autoantigenic fragments: DNA-PK_{CS} from amino acids 2699 to 4096; DNA-PK_{CS} from amino acids 3211 to 4096; PARP from amino acid 1 to 537; PARP from amino acids 538 to 1004; NuMA from amino acids 412 to 2111 and NuMA from amino acids 1 to 1799.

An aspect of this invention is a method of treating a patient in need of treatment for a malignancy. In a preferred embodiment, at least one enzyme of a lymphocyte granule is contacted with the malignant cells from the patient. This can produce a mixture containing autoantigenic fragments derived from the malignant cells. The fragments are administered to the patient, preferably with an adjuvant, to stimulate an immune response against the malignant cells.

An aspect of this invention is an assay for the detection of an autoantigenic fragment in a patient. In one embodiment, the presence or absence of the fragment in a patient sample is an indication of the presence or absence of an autoimmune condition in the patient. In a preferred embodiment, a sample from the patient is contacted with an antibody that specifically binds to a cryptic epitope of an autoantigenic fragment. Preferably, the fragment has at least one terminus derived from the cleavage of an autoantigen by granzyme B at a site that is not cleaved by a caspase. The presence or absence of the binding of the antibody to the autoantigenic fragment is then assessed as an indication of the presence or absence of an autoimmune condition in a patient. In an alternative embodiment, the detection of an antibody that binds an autoantigenic fragment is an indication of the presence or absence of an autoimmune condition in the patient. In this embodiment a sample from the patient is contacted with an autoantigenic fragment having at

least one terminus derived from cleavage by a granule enzyme. Detection of the presence or absence of the binding of an antibody in the sample to the autoantigenic fragment is an indication of the presence or absence of an autoimmune condition in the patient.

5 An aspect of this invention is a method of making an autoantigenic fragment from an autoantigen. In a preferred embodiment, one isolates cells containing at least one autoantigen and contacts the cells with a lymphocyte granule enzyme to produce a mixture containing at least one autoantigenic fragment. In a further
10 embodiment one isolates at least one autoantigenic fragment from the mixture. In a preferred embodiment one purifies at least one autoantigen and contacts the purified autoantigen with granzyme B. In a further preferred embodiment, one purifies one or more of the following autoantigens for contacting with granzyme B: DNA-PK_{CS},
15 PARP and NuMA. In each embodiment the granule enzyme can be isolated from the granules of a lymphocyte, *e.g.*, a cytotoxic T lymphocyte (CTL), a natural killer cell (NK), a lymphokine activated killer cell (LAK) or cells of the YT cell line.

In all aspects of this invention, granzyme B can be used in
20 particular embodiments. the enzyme can be purified from the granules of granule containing lymphocytes or can be prepared by recombinant techniques.

Definitions:

25 As used herein, "treatment" includes the therapeutic or prophylactic application of a composition to a patient. A treatment can prevent, moderate or cure a disease in the patient. A disease is moderated in a patient when the treatment lessens the severity or frequency of at least one symptom associated with the disease. A
30 treatment can moderate a disease by: (1) prophylactic administration of a composition to a patient free of a disease to lessen the impact of at least one symptom of the disease when it does occur or (2) therapeutic administration to a patient having a disease to lessen at least one symptom of the disease.

As used herein, a "patient" is an animal, particularly including a human.

As used herein, an "autoimmune condition" is the presence of, or the predisposition for the development of, an autoimmune disease or an autoimmune response in a patient.

As used herein, an "autoantigen" is a cellular molecule and usually is a protein. An autoantigen is typically not antigenic because the immune system is tolerized to its presence in the body under normal conditions. An autoantigen will typically include at least one cryptic epitope. An autoantigen can be produced by natural cells, using recombinant methods, or through chemical synthesis, as appropriate.

As used herein, an "autoantigenic fragment" is a degradation product of an autoantigen. An autoantigenic fragment is antigenic because the immune system is not tolerized to its presence in the body. Autoantigenic fragments usually display cryptic epitopes to the immune system. An autoantigenic fragment can be produced by natural cells, through the action of at least one granule enzyme in a cellular or cell free system, using recombinant methods, or through chemical synthesis, as appropriate.

As used herein, an "autoantibody" is an antibody produced by the immune system of an animal in response to the present of an autoantigenic fragment. An autoantibody can bind to the autoantigenic fragment, the autoantigen from which the fragment is derived, or both.

As used herein, a "granule containing lymphocyte" is meant to include all lymphocytes that contain granules. In particular, the term is used to include the family of cell types sometimes referred to as cytotoxic lymphocytes, to include cell lines derived from these cells and to include cytotoxic lymphocyte-like cell lines, preferably the YT cell line. Preferred cells are the granule containing lymphocytes known in the art as cytotoxic T lymphocytes (CTL), natural killer cells (NK) and lymphokine activated killer cells (LAK).

As used herein, a "lymphocyte granule enzyme" or "granule enzyme" is an enzyme that is found in the granules of a granule containing lymphocyte. A granule enzyme can be purified from a lymphocyte granule by methods commonly employed in the art of

protein purification. Additionally, a granule enzyme can be prepared by cloning the gene for the enzyme and the enzyme then is prepared using methods commonly used in the production of recombinant enzymes.

As used herein, "purified" and/or "isolated" are used interchangeably to stand for the proposition that the protein(s) and polypeptide(s), or respective fragment(s) thereof in question has been removed from its *in vivo* environment. A protein or fragment thereof is considered "purified" and/or "isolated" when it is obtained at a concentration at least about five-fold to ten-fold higher than that found in nature. A protein or fragment thereof is considered substantially pure if it is obtained at a concentration of at least about 100-fold higher than that found in nature. A protein or fragment thereof is considered essentially pure if it is obtained at a concentration of at least about 1000-fold higher than that found in nature. A protein is sometimes referred to as partly purified if it is at least purified or isolated but it is not essentially pure. A chemically synthesized protein is considered to be substantially purified when purified from its chemical precursors. A purified or isolated protein can be manipulated by the skilled artisan, such as but not limited to obtaining the protein or protein fragment in quantities that afford the opportunity to generate polyclonal antibodies, monoclonal antibodies, amino acid sequencing, and peptide digestion. Therefore, the autoantigenic fragments claimed herein can be present in cell lysates or in a substantially or essentially pure form.

25 Abbreviations:

Ac-DEVD-CHO, N-(N-Ac-Asp-Glu-Val)-3-amino-4-oxobutanoic acid; Ac-YVAD-CHO, N-(N-Ac-Tyr-Val-Ala)-3-amino-4-oxobutanoic acid; CTL, cytotoxic T lymphocytes; DNA-PK_{cs}, DNA-dependent protein kinase catalytic subunit; LAK cells, lymphokine-activated killer cells; NK cells, natural killer cells; NuMA, nuclear mitotic apparatus protein; PARP, poly(ADP-ribose)polymerase; PI, propidium iodide

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides autoantigenic fragments and methods for their use in the treatment of autoimmune disease. Also provided are assays for detecting an autoimmune condition in an animal, including the presence of an autoimmune disease.

The present invention makes use of the discovery that a class of previously unrecognized autoantigenic fragments is generated during the form of apoptosis triggered by the action of the contents of lymphocyte granules on cells. In particular, enzymes found in lymphocyte granules are discovered to cleave proteinaceous cellular autoantigens to yield previously unrecognized autoantigenic fragments. Granzyme B is found to be an important granule enzyme in this generation of autoantigenic fragments. This enzyme was previously known to cleave some of the pro-caspase enzymes to yield active caspases. Granzyme B is now found to directly cleave certain of the substrates of the caspases at different sites to produce novel autoantigenic fragments.

The present invention provides a method of producing granzyme generated autoantigenic fragments *in vitro* in several ways. In a preferred embodiment, purified granzyme B can be contacted with purified substrates to produce the autoantigenic fragments. Partly or wholly purified enzyme can also be used on partially or wholly purified substrates or cellular lysates containing the substrates. In this case, the autoantigenic fragments can be purified after cleavage of the substrates. Additionally, the contents of granules can be used to produce autoantigenic fragments of cellular components, including the autoantigenic fragments of caspase substrates created by the action of granzyme B, by application of granule contents, or purified substrates or tissue samples isolated from a patient. In some embodiments, the cells of the tissue can be disrupted by lysis or mechanical breakage to release the contents of the cells before contacting the cells with the contents of the granules.

The autoantigenic fragments provided herein can be used in a treatment to tolerize a patient to the presence of the autoantigenic fragments. Once tolerized, the patient would not develop an

autoimmune disease associated with the later appearance of the fragments in the patient. Tolerizing strategies involve purification of relevant autoantigenic fragments in a non-aggregated form. Low doses of the fragments are injected in pharmaceutically acceptable carriers, preferably without an adjuvant to induce low-zone tolerance.

The present invention also provides a method of generating an autoimmune response against certain cells in a patient. For example, if it is desirable to generate an autoimmune response against malignant cells in a patient, one can isolate a sample of the cells from the patient and contact the cells with the contents of granules isolated from granule containing lymphocytes. The action of the granule contents on the cells can produce autoantigenic fragments therefrom. In some embodiments, the malignant cells can be disrupted by lysis or mechanical breakage to release the contents of the cells before contacting the cells with the contents of the granules. In any case, the resulting mixture or purified components therefrom can be administered to the patient. Because the autoantigenic fragments produced in this way are the same as those produced *in vivo* by the action of granule containing lymphocytes, including *e.g.*, CTLs, NK and LAK cells, on the malignant cells, the immune system is thereby stimulated to generate a response against the malignant cells. Therefore, the present invention provides a method to heighten or stimulate the natural immune system processes to act against particular types of cells such as malignant cells.

The autoantigenic fragments useful in the invention described herein display cryptic epitopes. These epitopes are revealed to the immune system after cleavage of the precursor protein by the enzymes contained in granules to yield the autoantigenic fragments. There is a persuasive body of literature that reports that the highly specific humoral immune response to autoantigens in autoimmune disease is T cell-dependent, and that flares in autoimmune disease result when this primed immune system is rechallenged with a self-antigen (reviewed in Burlingame, R.W., *et al.*, 1993; Diamond, B., *et al.*, 1992; Radic, M.Z. and M. Weigert. 1994). However, the mechanisms responsible for initiation of the primary immune response to these molecules, and for subsequently stimulating the secondary response to

targeted antigens, have not been completely elucidated (Bach, J.F. and S. Koutouzov. 1997; Sercarz, E.E. and S.K. Datta. 1994). Several studies report that a potential for T cell autoreactivity resides in the immunological non-equivalency of different areas of self-molecules, since tolerance is only induced to dominant determinants which are generated and presented at suprathreshold concentrations during natural processing of whole protein antigens (reviewed in (Sercarz, E.E., *et al.*, 1993; Lanzavecchia, A. 1995). Those determinants which are not generated at all, or are generated at subthreshold levels during antigen processing (termed cryptic), do not tolerize T cells. Thus, potentially autoreactive T cells recognizing this cryptic self epitope are allowed to persist.

The mechanisms through which autoimmunity can arise when normally cryptic determinants become visible to the immune system has received increased attention. Several experimental systems have now provided evidence that the balance of dominant versus cryptic epitopes in a self molecule can be profoundly influenced by forces which alter the 'immunological' structure of molecules (Lanzavecchia, A. 1995). Examples include the revelation of cryptic epitopes through novel cleavage (Bockenstedt, L.K., *et al.*, 1995; Mamula, M.J. 1993), or through altered conformation induced by high affinity ligand binding (e.g. to an antibody or receptor molecule (Salemi, S., *et al.*, 1995; Simitsek, P.D., *et al.*, 1995; Watts, C. and A. Lanzavecchia. 1993). The unique, high-titer autoantibody responses that characterize different autoimmune diseases can therefore be viewed as the immunologic impression of the initiating events that revealed suprathreshold concentrations of non-tolerized structure in a pro-immune context, thus satisfying the stringent criteria for initiation of a primary immune response (Casciola-Rosen, L. and A. Rosen. 1997).

The alteration of the structure of autoantigens during apoptosis is an important feature which underlies targeting of specific molecules by the immune system. Understanding whether and how the structure of those autoantigens are cleaved during apoptosis, provides insights into the role of apoptosis in initiation of the autoimmune response. One such mechanism involves the action of the enzymes in

contents of granules found in granule containing lymphocytes and is described herein. The contents of these granules are shown to act to reveal cryptic epitopes by cleaving autoantigens to autoantigenic fragments. In a preferred method, granzyme B cleaves autoantigenic proteins to create preferred autoantigenic fragments that display cryptic epitopes to the immune system.

Recent reports using caspase inhibitors have emphasized the contribution of caspase-independent pathway(s) when target cell death is induced by cytotoxic lymphocyte granule exocytosis (Sarin *et al.*, 1997; Talanian *et al.*, 1997). The results presented here demonstrate that several of the downstream substrates cleaved by the caspase family of proteases during apoptosis are also directly and efficiently cleaved by granzyme B both *in vitro* and in target cells undergoing lymphocyte granule-induced cytotoxicity. This confirms the existence of efficient caspase-independent proteolytic pathways during this form of cell death.

Not all downstream substrates of caspase-3 are cleaved by granzyme B with similar efficiency. In the case of PARP and U1-70kDa, cleavage efficiency by caspase-3 exceeds that by granzyme B by more than 200 fold, while NuMA and DNA-PK_{CS} are cleaved with similar efficiencies by granzyme B and caspase-3 (Table I). The relative cleavage efficiency of a given substrate by the two proteases appears to account for the results observed when caspase-3 activity is inhibited in intact cells. Thus, while PARP cleavage during granule-mediated cytotoxicity is inhibited by Ac-DEVD-CHO, cleavage of DNA-PK_{CS} and NuMA are only minimally affected. It is therefore believed that caspase-independent components of granule-mediated death are generated through altering the function of those downstream substrates that are efficiently cleaved by granzyme B (and potentially other granule proteases).

The results presented herein are consistent with recent studies in which granzyme B-induced proteolysis of PARP and U1-70kDa (both inefficient direct substrates for granzyme B) in an intact cell model was well inhibited by caspase inhibitors (Talanian *et al.*, 1997). The results differ from a previous study which failed to demonstrate unique DNA-PK_{CS} fragments during CTL-induced target cell death, or by granzyme B *in vitro* (Song *et al.*, 1996a). Different antibodies to DNA-

PK_{CS} used in the two studies may account for the failure of the previous study to find that a unique 100kDa C-terminal fragment is generated by granzyme B.

The results indicate that direct and efficient cleavage of NuMA and DNA-PK_{CS} by granzyme B achieves functional effects similar to those caused by caspase cleavage during other forms of apoptosis. This is supported by the demonstration that the cleavage of DNA-PK_{CS} by granzyme B completely abrogates its kinase activity (FIG. 3). This granzyme B-mediated cleavage of DNA-PK_{CS} differs from caspase-3-mediated cleavage of this substrate in several ways: (i) The presence of DNA ends renders the granzyme B-mediated cleavage significantly less efficient, while DNA ends are required for efficient cleavage of DNA-PK_{CS} by caspase-3 (Casciola-Rosen *et al.*, 1996; Song *et al.*, 1996b) (ii) While caspase-3 cleavage of DNA-PK_{CS} decreases its kinase activity to approximately 60% of control levels (Casciola-Rosen *et al.*, 1996), cleavage by granzyme B completely abrogates kinase activity. These data indicate that granzyme B cleaves and fully inactivates DNA-PK_{CS} very early during granule-mediated cytotoxicity, before significant internucleosomal DNA cleavage has occurred, and thus before caspase-3 can efficiently cleave DNA-PK_{CS}. Once abundant DNA ends have been generated, cleavage by caspase-3 should predominate (with the resulting fragments possessing residual kinase activity).

Although the functional consequences of substrate cleavage by caspases during apoptosis *in vivo* is not known for most substrates, recent reports demonstrate that cleavage can activate critical pro-apoptotic activities (e.g. activation of DNA fragmentation factor by caspase-3 (Liu *et al.*, 1997)). In other cases, reports suggest that substrate cleavage might disable important structural and homeostatic functions (Casciola-Rosen *et al.*, 1996; Ghayur *et al.*, 1996). It is therefore reported that the ability of cytolytic lymphocyte granule proteases to directly cleave some caspase substrates underlies the caspase-independent component of the death pathway induced by these cells (Sarin *et al.*, 1997). The ability of the lymphocyte granule-induced

cytotoxicity pathway to generate a novel form of nuclear condensation even in the absence of caspase activity, is consistent with this theory.

Studies on macromolecular and tetrapeptide substrates provide insights into the substrate specificity of granzyme B. For example, the direct cleavage of PARP by granzyme B occurs at VGPD537-S538 (Froelich *et al.*, 1996a), while processing of several caspase precursors occurs at IETD sites (Fernandes-Alnemri *et al.*, 1996; Ramage *et al.*, 1995; Srinivasula *et al.*, 1996; Yamin *et al.*, 1996). The studies presented here demonstrate that granzyme B cleaves a macromolecular substrate (DNA-PK_{CS}) at VGPD-F2699, a site consistent with the tetrapeptide substrate specificity of granzyme B defined using a combinatorial tetrapeptide library (Thornberry *et al.*, 1997). Interestingly, granzyme B also directly cleaves this substrate at a nearby DEVD-N2713, suggesting that the determinants of macromolecular substrate specificity are more complex than those contained in the tetrapeptides tested.

Using the fragment sizes of NuMA and DNA-PK_{CS} generated by granzyme B, the known epitope specificity of several of the antibodies used, and the known substrate specificity of granzyme B, these studies indicate that the most likely granzyme B cleavage site generating the 100kDa C-terminal fragment of DNA-PK_{CS} is VDQD3210-G3211. Similarly, the most likely granzyme B cleavage site in NuMA occurs at VLGD411-V412. Several studies have addressed the cell biology of granzyme B during perforin/granzyme B-induced apoptosis (Froelich *et al.*, 1996b; Shi *et al.*, 1997; Trapani *et al.*, 1996). These studies reported that granzyme B autonomously enters the cytoplasm of target cells. That event alone does not induce target cell apoptosis. In the presence of perforin, however, apoptosis is induced in target cells. That event is accompanied by the rapid enrichment of granzyme B in nuclei and nucleoli of target cells (Jans *et al.*, 1996; Pinkoski *et al.*, 1996; Trapani *et al.*, 1996). It is therefore of importance that the granzyme B substrates described here are nuclear proteins. These substrates function in both structural and homeostatic pathways which are impacted by the caspases during apoptosis. The observation that a component of granule-induced cell death is caspase-independent, taken

together with the ability of caspases and granzyme B to efficiently cleave a common subset of downstream substrates at different sites during granule-induced cytotoxicity, highlights the importance of proteolysis of those common substrates in generating the apoptotic phenotype.

5 Table II).

In the light of several descriptions of viral or endogenous caspase inhibitors (Beidler *et al.*, 1995; Bump *et al.*, 1995; Irmeler *et al.*, 1997; Thome *et al.*, 1997; Xue and Horvitz, 1995), as well as long-lived cells or tumor cells which express low levels of specific caspase family members (Krajewska *et al.*, 1997; Krajewski *et al.*, 1997), the caspase-independent activity of granzyme B provides the host with apoptotic effector mechanisms that are insensitive to inhibitors of the signaling or execution components of the apoptotic cascade.

Many of the downstream caspase substrates described to date are autoantigens targeted in human systemic autoimmune diseases (Casciola-Rosen *et al.*, 1995; Casciola-Rosen *et al.*, 1994; Casciola-Rosen *et al.*, 1996; Casiano *et al.*, 1996; Greidinger *et al.*, 1996). The present demonstration that several of these substrates are also directly cleaved by granzyme B, generating unique fragments not generated during any other form of apoptosis studied to date, demonstrates how non-tolerized determinants of autoantigens can be revealed during certain forms of CTL-mediated apoptotic death.

In the experimental results presented herein, it is demonstrated that DNA-PK_{CS} and NuMA are directly cleaved by granzyme B, both *in vitro* and in cells undergoing granule-induced cytotoxicity. Although the efficiency of cleavage of these substrates is similar to those observed for caspase 3-mediated cleavage, the fragments generated by the 2 proteases are distinct. Since caspases appear to initiate apoptosis by altering the function of downstream substrates (either by decreasing the function of the intact substrate, or by generating fragment(s) with pro-apoptotic activity), it is believed that direct cleavage of caspase substrates by granzyme B during cytotoxic lymphocyte granule-induced apoptosis plays an important role in caspase-independent target cell death. The ability of the contents of cytotoxic lymphocyte granules to bypass the requirement for caspases in

the death pathway may guarantee the demise of target cells whose caspase pathway is incomplete or under strict endogenous or exogenous regulatory control.

The action of granzyme B in producing particular autoantigenic fragments from particular autoantigens is exemplified herein. However, the general understanding of the role of granule proteases as described and exemplified in the granzyme B model system allows one to generate these and other autoantigenic fragments. The autoantigenic fragments produced can be used in the preparation of pharmaceutical compositions, for treating patients at risk for or suffering from autoimmune diseases and cancer, and in assays for assessing the presence or absence of an autoimmune condition in a patient.

The following examples are presented by the way of illustration and, because various other embodiments will be apparent to those in the art, the following examples are not to be construed as a limitation on the scope of the invention.

EXAMPLE 1

General Materials and Methods

Materials. Purified DNA-dependent protein kinase (DNA-PK) and SP1 were purchased from Promega (Madison, WI). ATP was purchased from Fluka (Ronkonkoma, NY), and ^{32}P -ATP was from Du Pont/NEN (Wilmington, DE). Ac-DEVD-CHO and Ac-YVAD-CHO were manufactured by Merck (Rahway, NJ). Caspase-3 was purified as described (Nicholson *et al.*, 1995). Patient sera were used to immunoblot the nuclear mitotic apparatus protein (NuMA), poly(ADP-ribose) polymerase (PARP) and DNA-PK_{CS} (Casciola-Rosen *et al.*, 1995; Greidinger *et al.*, 1996). Monoclonal antibodies can be made by methods known in the art. Two different monoclonal antibodies, designated 18-2 and 25-4 (kind gifts from Dr. Tim Carter, St. Johns University, Jamaica, NY) were also used to detect DNA-PK_{CS} by immunoblotting (see Table II

for a summary of the antibodies used to detect DNA-PK_{CS} and its cleaved fragments). Rabbit polyclonal antibodies to caspases were raised against the large subunits of caspase-3 and caspase-7, using methods commonly known in the art. Immunoblotted proteins were detected using the
 5 SUPERSIGNALTM substrate system (Pierce, Rockford, IL), according to the manufacturer's instructions.

In vitro cleavage of purified DNA-PK_{CS} and

[³⁵S]methionine-labeled caspase-3 precursor, caspase-7 precursor, PARP, and NuMA. cDNAs for caspase-3, caspase-7, NuMA and PARP
 10 were used to drive the synthesis of [³⁵S]methionine-labeled proteins by coupled transcription/translation in rabbit reticulocyte lysates. For all purified substrates, cleavage reactions were performed in buffer consisting of 50 mM Hepes pH 7.4, 10% sucrose and 5 mM DTT in the presence of the granzyme B concentrations indicated in FIG. 1. After
 15 incubation at 37⁰C for 15 min, reactions were terminated and samples were electrophoresed on 10% (DNA-PK_{CS}, NuMA), 12% (PARP) or 15% (caspases 3 and 7) SDS-polyacrylamide gels. Radiolabeled proteins and their fragments were visualized by fluorography. Intact and cleaved DNA-PK_{CS} were visualized by immunoblotting with monoclonal antibody
 20 18-2 (Casciola-Rosen *et al.*, 1995).

Calculation of catalytic constant values. Catalytic constant (k_{cat}/k_m) values were calculated essentially as described (Casciola-Rosen *et al.*, 1996). Briefly, subsaturating substrate concentrations were used in each *in vitro* reaction, and product appearance was assumed to
 25 be a first order process. Substrate and product bands on autoradiograms were scanned by densitometry. Several appropriate densitometry systems are available, *e.g.* PDI Discovery System, with Quantity One Software, Protein Databases, Inc., (Huntington Station, NY). k_{cat}/K_m values were calculated by fitting the dose-response data to the first order
 30 rate equation: percent substrate cleavage = 100*(1-e(-(k_{cat}*[E]/K_m)*time)).

In vitro cleavage of endogenous DNA-PK_{CS}, NuMA, and PARP in HeLa lysates. Control HeLa lysates were prepared using methods commonly applied in the art, as described in Casciola-Rosen *et al.*, 1994. 12.5 nM purified granzyme B or 105 pM purified caspase-3

were then added to the lysates, in the absence or presence of 100nM Ac-DEVD-CHO or Ac-YVAD-CHO. The mixtures were incubated for 15 min on ice to facilitate binding of the inhibitors to proteases prior to performing cleavage reactions at 37°C for 60 min. After

- 5 electrophoresing the samples on 10% SDS-polyacrylamide gels containing 0.087% bisacrylamide, the intact proteins and their cleaved fragments were visualized by immunoblotting.

- Determination of granzyme B cleavage sites in DNA-PK_{CS} by P1 Asp mutagenesis.* A partial cDNA clone for DNA-PK_{CS}, encoding
10 Met²⁵⁶⁶ through Leu²⁹²⁸ (the region containing the caspase-3 cleavage site as well as the putative granzyme B site), was amplified by reverse-transcriptase PCR from HeLa cell poly(A)⁺ RNA using primers containing 5' EcoRI and 3' XbaI restriction enzyme-adapters. After ligation into the corresponding restriction sites of pBluescript II SK⁺
15 (Stratagene), this clone was used as template for mutagenesis by overlap-extension PCR to generate clones containing D2698A (P1 of the putative granzyme B site) and D2712A (P1 of the known caspase-3 site) modifications. [³⁵S]-Radiolabeled polypeptides were generated by coupled *in vitro* transcription/ translation, and then incubated with
20 either recombinant caspase-3 (8 nM) or purified YT cell-derived granzyme B (8 nM) for 60 min at 37°C in a buffer composed of 50 mM Hepes/KOH (pH 7.0), 10% (w/v) sucrose, 2 mM EDTA, 0.1% (w/v) CHAPS, 5 mM dithiothreitol. The resulting cleavage products were resolved on SDS-polyacrylamide gels (10-20% gradient gels) and
25 visualized by fluorography.

- In vivo cleavage of endogenous DNA-PK_{CS}, NuMA and PARP during YT cell granule content-induced cytotoxicity of intact K562 cells.* Intact cytoplasmic granules were purified from YT cells, and the granule contents were isolated using known methods (Tschopp, 1994).
30 Some of these preparations were used for the further purification of granzyme B (Tschopp, 1994). Purity of the protease was confirmed by silver staining of overloaded SDS-polyacrylamide gels. The cytotoxic effects of YT granule contents were determined as follows: Jurkat T cells or K562 cells were radiolabeled with 100μCi/ml [⁵¹Cr] sodium chromate,

washed, resuspended in Ca^{2+} -free HBSS, and then incubated with granule contents (0-5 μl) and 1mM CaCl_2 for increasing times. The percentage specific [^{51}Cr] was calculated using the following formula:

- 5 % specific lysis = [(sample cpm- spontaneous cpm)/(maximum cpm – spontaneous cpm)] x 100. *In vivo* experiments to assay the effect of YT cell granule contents on DNA-PK CS , NuMA and PARP in intact K562 cells were performed as follows: K562 cells were washed twice with PD (2.7mM KCl, 1.5mM KH_2PO_4 , 137mM NaCl, 8mM Na_2HPO_4), then
 10 resuspended at 1.7×10^7 cells/ml in PD in the absence or presence of 100 μM Ac-DEVD-CHO, and incubated at 37°C for 30 min. Aliquots containing 3.4×10^5 cells were incubated for a further 90 min at 37°C with 1mM EDTA or 1mM Ca^{2+} and 2 μl of YT granule contents (which induces 20-40% specific chromium release in 60 min, and characteristic internucleosomal DNA degradation. The total reaction volume of each
 15 sample was 30 μl . The reactions were terminated by boiling in SDS gel buffer and samples were electrophoresed and immunoblotted as described above. Experiments were performed using 2 different preparations of granule contents and 4 different cell types (K562, Jurkat, HeLa, primary human myoblasts).

- 20 *Confocal Immunofluorescence microscopy.* Morphologic experiments were performed on HeLa cells grown on No. 1 glass coverslips. Coverslips were washed three times with ice-cold HBSS without Ca^{2+} , prior to incubation (4°C, 30 min) with 25 μl of HBSS minus Ca^{2+} containing 0.8 μl of YT cell granule contents (see above), in the
 25 presence or absence of 200 μM Ac-DEVD-CHO. 25 μl of HBSS containing 2mM CaCl_2 was then added to each coverslip (mixed well by repeated, gentle aspiration), followed by incubation in a humidified chamber at 37°C for 60 min. The cells were then fixed in 4% paraformaldehyde (4°C, 5 min), permeabilized with acetone (4°C, 15 sec), and stained
 30 sequentially with antibodies to PARP or NuMA, propidium iodide and DAPI as described (Casciola-Rosen *et al.*, 1994a). Coverslips were mounted on glass slides with Permafluor (Lipshaw, Pittsburgh, PA), and confocal microscopy was performed on a scanning confocal microscopy system (LSM 410, Carl Zeiss, Inc., Thornwood, NJ).

LAK cell-mediated cytotoxicity. LAK cells were obtained by culturing human PBMCs for 4 days in LAK medium (RPMI supplemented with 10mM Hepes pH 7.4, L-glutamine, 2% autologous plasma), and 1000 Cetus units/ml of hrIL-2 (Chiron Therapeutics, Emeryville, CA) (Topalian *et al.*, 1989). Fas-negative target cells (K562 erythroleukemia cells) were resuspended at 1.3×10^6 cells/ml in LAK medium in the presence or absence of 100 μ M Ac-DEVD-CHO, and incubated at 37°C for 60 min, prior to co-incubation with LAK effector cells (effector:target ratio of 5:1) for 4 h. After 2 washes with PD, cells were lysed and boiled in SDS sample buffer, and PARP, DNA-PK_{CS} and NuMA were assayed by immunoblotting as described above.

Kinase assay. Kinase assays were performed on intact DNA-PK_{CS} or DNA-PK_{CS} that had first been cleaved by granzyme B as follows. Reaction mixtures containing 10 mM Hepes pH 7.4, 2 mM MgCl₂, 10 mM KCl, 2.7 mM DTT, and 50 ng DNA-PK_{CS} in the absence or presence of 12.5 nM purified granzyme B, were incubated for 13.5 min at 37°C. Kinase reactions were subsequently initiated by adding 100 ng SP1 and 150 μ M ATP containing 1.5 μ Ci [³²P]-ATP (3000Ci/mmol), in the absence or presence of 10 μ g/ml sheared herring sperm DNA (Promega). Samples were incubated at 37°C for 10 min (well within the linear range of the assay, data not shown), before terminating the reactions by adding SDS gel buffer and boiling. After electrophoresing the samples on 8% SDS-PAGE, SP1 phosphorylation was detected by autoradiography, and quantitated by densitometry. Cleaved status of the kinase was confirmed in parallel by immunoblotting.

EXAMPLE 2

DNA-PK_{CS} and NuMA are very efficient substrates for purified granzyme B.

Granzyme B has previously been reported to cleave the precursors of several caspases (including caspases 3, 7 and 10), resulting in activation of their proteolytic activity. The catalytic efficiency of cleavage of these substrates by granzyme B serves as a useful

standard against which granzyme B-mediated cleavages of other substrates can be compared. Purified [^{35}S]methionine-labeled precursors of caspase-3 and caspase-7, or THP.1 cytosols (containing these precursor proteases) were incubated *in vitro* with increasing concentrations of purified granzyme B. The dose-response data obtained (FIG. 1) was used to calculate catalytic constant ($k_{\text{cat}}/K_{\text{M}}$) values of $1.8 \pm 0.6 \times 10^5 \text{M}^{-1}\text{s}^{-1}$ (radiolabeled substrate) and $1.9 \pm 0.1 \times 10^5 \text{M}^{-1}\text{s}^{-1}$ (immunoblotting) for caspase-7, and $3.6 \pm 1.0 \times 10^4 \text{M}^{-1}\text{s}^{-1}$ (radiolabeled substrate) and $2.3 \pm 0.4 \times 10^4 \text{M}^{-1}\text{s}^{-1}$ (immunoblotting) for caspase-3 (Table I). Thus, granzyme B cleaves caspase-7 approximately 6 fold more efficiently than caspase-3, consistent with previous reports (Talanian *et al.*, 1997).

It was then determined whether any of the downstream substrates known to be cleaved by caspases during apoptosis were also directly cleaved by granzyme B, and the efficiency of cleavage of each substrate by the two proteases was compared. Purified DNA-PK $_{\text{CS}}$ was very efficiently cleaved by granzyme B in the absence of added DNA ($k_{\text{cat}}/K_{\text{M}} = 2.5 \pm 0.8 \times 10^6 \text{M}^{-1}\text{s}^{-1}$, see FIG. 1 and Table I), making it the best substrate for granzyme B described to date, with a cleavage efficiency two orders of magnitude better than that described for granzyme B-mediated cleavage of the caspase-3 precursor. When similar experiments were performed in the presence of $10 \mu\text{g/ml}$ DNA, DNA-PK $_{\text{CS}}$ cleavage was decreased by approximately 90% .

Caspase-3-mediated cleavage of DNA-PK $_{\text{CS}}$ was also extremely efficient ($k_{\text{cat}}/K_{\text{M}}$ value = $7.5 \pm 0.8 \times 10^6 \text{M}^{-1}\text{s}^{-1}$). In this case, efficient cleavage was only obtained in the presence of DNA ((Casciola-Rosen *et al.*, 1995), and Table I). NuMA, a nuclear matrix protein that is cleaved in apoptotic cells by an unidentified protease with features of the caspase family, was also very efficiently cleaved by granzyme B ($k_{\text{cat}}/K_{\text{M}}$ value = $5.4 \pm 1.4 \times 10^5 \text{M}^{-1}\text{s}^{-1}$, see Table I). The efficiency of this cleavage was one order of magnitude greater than that observed for granzyme B-mediated processing of the caspase-3 precursor. NuMA was also efficiently cleaved by purified caspase-3, with a $k_{\text{cat}}/K_{\text{M}}$ value of = $5.0 \pm 1.0 \times 10^5 \text{M}^{-1}\text{s}^{-1}$. (Table I). In contrast to DNA-PK $_{\text{CS}}$ and NuMA (where granzyme B- and caspase-3-mediated cleavages are similarly efficient),

PARP was a relatively poor substrate for granzyme B, with a k_{cat}/K_m value ($2.3 \pm 1.8 \times 10^4 \text{ M}^{-1}\text{s}^{-1}$) that is approximately 200 fold lower than that for caspase-3 (see Table I). Granzyme B was also a poor catalyst for cleaving U1-70kDa, with k_{cat}/K_m values $<10^3 \text{ M}^{-1}\text{s}^{-1}$. The efficiency of substrate cleavage by granzyme B is therefore similar to caspase-3 for some substrates (*e.g.* DNA-PK_{CS} and NuMA), while it is more than 2 orders of magnitude less efficient for others (*e.g.* PARP and U1-70kDa).

EXAMPLE 3

10 Different substrate fragments are detected after cleaving autoantigens *in vitro* with granzyme B or caspase-3.

To directly compare the fragments generated by granzyme B and caspase-3, purified DNA-PK_{CS}, *in vitro* translated [35S]methionine-labeled PARP, and endogenous substrates (NuMA and DNA-PK_{CS} in HeLa cell lysates) were incubated with protease and electrophoresed in adjacent lanes. When granzyme B was used to cleave DNA-PK_{CS}, fragments of 100kDa and 250kDa were generated, (detected by immunoblotting using antibodies recognizing the C-terminus or N-terminus of DNA-PK_{CS}, respectively) (FIG. 2, lanes 2 & 5; and Table II). In contrast, caspase-3 cleavage yielded a 150kDa C-terminal fragment (FIG. 2, lane 3) and a 250kDa N-terminal fragment (FIG. 2, lane 6).

Granzyme B-mediated cleavage of NuMA generated a novel fragment migrating at 175kDa on SDS-PAGE, which was distinct from the 185kDa fragment detected after cleavage with caspase-3 (FIG. 2, lanes 7-9). Similarly, novel fragments of PARP migrating at 72, 62 and 42kDa were detected after incubation with granzyme B; these differed from the 89 and 24 kDa fragments generated by caspase-3-mediated cleavage of PARP (FIG. 2, lanes 10-12 and Table II). Granzyme B therefore directly cleaves several of the downstream substrates of caspase-3 *in vitro*. In all cases, the fragments generated by granzyme B differ from those generated by caspase-3.

EXAMPLE 4

Kinase activity of DNA-PK_{CS} is abolished by granzyme B cleavage.

To determine the effect of granzyme B-mediated cleavage on the kinase activity of DNA-PK_{CS}, the ability of intact and cleaved DNA-PK_{CS} to phosphorylate the SP1 transcription factor was quantitated. The kinase activity of DNA-PK_{CS} was entirely DNA-dependent (FIG. 3, lanes 1 and 2). When DNA-PK_{CS} was pretreated with 12.5nM granzyme B for 13.5 min at 37°C (which cleaves DNA-PK_{CS} (FIG. 1)), followed by the addition of DNA, [³²P]ATP and SP1 (which initiate the phosphorylation reaction), kinase activity was entirely abolished (FIG. 3, lane 3). Whether any of the fragments generated by granzyme B-mediated cleavage of DNA-PK_{CS} have novel activity(ies) in addition to the autoantigenic reactions demonstrated herein remains to be determined.

EXAMPLE 5

Granzyme B induces novel fragments of DNA-PK_{CS} and NuMA in cell lysates.

Previous studies reported that the cleavage of PARP that occurs in lysates of COS cells expressing granzyme B can be almost completely inhibited by the caspase inhibitor, Ac-DEVD-CHO (Darmon *et al.*, 1995). Because this compound is not an inhibitor of granzyme B, this observation suggests that this cleavage is mediated solely by caspases which have been activated by granzyme B in these extracts. The observation that PARP itself is cleaved by granzyme B approximately 200 fold less efficiently than by caspase-3 further supports this conclusion (see Table I). Since NuMA and DNA-PK_{CS} are cleaved with similar efficiencies by caspase-3 and granzyme B, it was of interest to determine whether these substrates are cleaved directly by granzyme B in cell lysates that contain caspase precursors. Therefore, an *in vitro* assay system was established in which cleavage of endogenous DNA-PK_{CS}, NuMA or PARP was monitored after addition of purified caspase-3 or

granzyme B. The assay was conducted in the presence or absence of the caspase inhibitor, Ac-DEVD-CHO.

Incubation of control extracts at 37°C for 60 minutes resulted in no significant cleavage of DNA-PK_{CS} or PARP, but was

- 5 associated with minimal cleavage of NuMA, leading to the production of minor fragments of 185 and 187kDa (FIG. 4, lane 1). When purified caspase-3 was added to control extracts, caspase-3-specific fragments of PARP (89kDa), NuMA (185kDa) and DNA-PK_{CS} (250kDa N-terminal fragment, 150kDa C-terminal fragment) were generated (FIG. 4, lane 2).
10 As demonstrated above, substrate fragments generated by the activity of caspase-3 are entirely abolished by 100nM Ac-DEVD-CHO (K_i caspase-1 = 17nM; K_i caspase-3 = 0.2nM; K_i caspase-6 = 31nM; K_i caspase-7 = 1nM; K_i caspase-8 = 0.92nM ; K_i caspase-9 = 60nM; K_i caspase-10 = 12nM) (FIG. 4, lane 3), but were unaffected by 100nM Ac-YVAD-CHO (K_i caspase-1 = 0.6nM; K_i caspase-3 and caspase-7 >10 μ M; K_i caspase-10 = 408nM) (FIG. 4, lane 4).

- In the presence of 12.5nM granzyme B, all three substrates were efficiently cleaved (FIG. 4, lane 5). The predominant PARP fragment induced by granzyme B co-migrated with the fragment
20 induced by caspase-3 (FIG. 4, lane 5). There was also a minor (<5%) fragment of 62kDa which corresponded with the fragment induced by granzyme B on purified substrate. In contrast, only a minor proportion of the NuMA fragments induced by granzyme B co-migrated with the fragment generated by caspase-3 (185kDa). The major, novel fragment
25 of 175kDa (FIG. 4, lane 5) corresponded with the fragment induced by granzyme B on purified *in vitro*-translated substrate. Granzyme B-mediated cleavage of DNA-PK_{CS} generated a 250kDa N-terminal fragment co-migrating with that generated by caspase-3 (FIG. 4, lanes 2 & 5), as well as a unique 100kDa C-terminal fragment which
30 corresponded with that induced by granzyme B on purified substrate (FIG.2, lane 2).

- When Ac-DEVD-CHO or 2mM iodoacetamide were added to the extracts 15 minutes prior to the addition of granzyme B, cleavage of PARP was almost entirely abolished (FIG. 4, lane 6). In contrast, the
35 cleavage of NuMA and DNA-PK_{CS} was only partially inhibited (FIG. 4,

lane 6) (20-40% inhibition in 5 separate experiments). Generation of caspase-3-specific fragments of both NuMA (185kDa) and DNA-PKcs (150kDa C-terminal fragment was abolished under these circumstances, while formation of the 250kDa N-terminal fragment of DNA-PKcs was inhibited by 20-40% (FIG. 4, lane 6).

Generation of granzyme B-specific fragments (175kDa NuMA fragment and 100kDa C-terminal DNA-PKcs fragment) was not affected by the inhibitors of caspase (FIG. 4, lane 6). The failure of Ac-DEVD-CHO to markedly inhibit generation of the 250kDa DNA-PKcs fragment indicates that the majority of DNA-PKcs cleavage detected after addition of granzyme B to lysates results from direct cleavage by this protease (rather than indirectly through activation of caspase-3).

Taken together, the results demonstrate that granzyme B competes with caspase-3 for cleavage of endogenous substrates in cell lysates. The outcome of this competition can be accurately predicted by comparing k_{cat}/K_m values: for those substrates where k_{cat}/K_m for cleavage by caspase-3 is greater than that of granzyme B (e.g. PARP), caspase-3 fragments are detected almost exclusively. For those substrates where the k_{cat}/K_m values are similar for the proteases, addition of granzyme B results in the formation of novel, granzyme B-specific fragments (e.g. DNA-PKcs and NuMA).

EXAMPLE 6

Granzyme B cleavage sites in DNA-PKcs

To address whether the 250kDa DNA-PKcs fragment generated by caspase-3 and granzyme B result from cleavage at the same and/or different, but closely-spaced sites, not distinguishable by SDS-PAGE, we generated a fragment of DNA-PKcs (Met²⁵⁶⁶ - Leu²⁹²⁸) encompassing both the known caspase-3 cleavage site at DEVD²⁷¹²-N²⁷¹³ (Casciola-Rosen *et al.*, 1996; Song *et al.*, 1996b), as well as the potential granzyme B cleavage site at VGPD²⁶⁹⁸-F²⁶⁹⁹. The relevant P1 aspartic acids (D²⁶⁹⁸ and D²⁷¹²) in this fragment were mutated to alanines (D²⁶⁹⁸A; D²⁷¹²A), and susceptibility of wild type and mutated

forms to cleavage by caspase-3 or granzyme B was assessed (FIG. 5). Cleavage of the wild type protein by caspase-3 resulted in two fragments of 26kDa and 20kDa (FIG. 5, lanes 1,2); this cleavage was entirely abolished by the D2712A mutation (FIG. 5, lanes 7,8), confirming that caspase-3 cleaves at DEVD2712-N2713. In addition to 20kDa and 26kDa fragments identical to those generated by caspase-3, granzyme B cleavage also resulted in fragments of 28kDa and 18kDa (FIG. 5, lane 3); these unique fragments were enhanced by the D2712A mutation (FIG. 5, lane 9), but were abolished by the D2698A mutation (FIG. 5, lane 6), placing a granzyme B cleavage site at VGPD2698-F2699. These data demonstrate that granzyme B can cleave at VGPD2698-F2699, a site predicted by previous studies using a combinatorial tetrapeptide substrate library (Thornberry *et al.*, 1997). Furthermore, granzyme B also cleaves at the caspase-3 cleavage site, DEVD2712-N2713. The C-terminal granzyme B-unique cleavage site is believed to be VDQD3210-G3211. This cleavage yields the approximately 100 kDa fragment from amino acids 3212 to 4096.

EXAMPLE 7

Granzyme B-specific fragments of DNA-PK_{cs} and NuMA are generated during cytotoxic lymphocyte granule-induced target cell death.

To determine whether the granzyme B-specific fragments demonstrated in cell lysates also occur when target cell death is induced by the contents of cytolytic lymphocyte granules, The consequences of granule-induced target cell death on substrate cleavage were examined. Granules containing perforin and granzymes were purified from the human NK cell line, YT. The granule contents were harvested and used to induce target cell cytotoxicity as described in Example 1. Several different target cells (Jurkat T cells, K562 erythroleukemia cells, human myoblasts or HeLa cells) were incubated with granule contents (~1.5 x 10⁷ YT cell equivalents/ml) in the presence of Ca²⁺. Rapid target cell lysis was induced (achieving ~20%-40% specific ⁵¹Cr release in 60 min). Target cell lysis by granule contents did not occur in the absence of Ca²⁺

(Podack and Konigsberg, 1984; Young *et al.*, 1986). DNA-PK_{CS}, NuMA and PARP were all cleaved rapidly after addition of granules, in a Ca²⁺-dependent manner (FIG. 6, lanes 2 & 3). As observed in the lysate system described above, cleavage of PARP resulted almost completely in the generation of the 89kDa caspase-3-specific fragment (FIG. 6, lane 3). Only small amounts of the 62kDa granzyme B-specific fragment were generated. Generation of the 89kDa fragment was completely inhibited by Ac-DEVD-CHO (FIG. 6, lane 4). In contrast, granule contents induced the formation of granzyme B-specific cleavage fragments of DNA-PK_{CS} and NuMA (FIG. 6, lane 3). The production of these fragments was not inhibited by Ac-DEVD-CHO (FIG. 6, lane 4).

Note that in the case of DNA-PK_{CS}, addition of granule contents also resulted in the formation of small amounts of 150kDa and 120kDa fragments, which were only well visualized after longer exposures of the X-ray film. Generation of these fragments was also entirely inhibited by Ac-DEVD-CHO.

EXAMPLE 8

Granzyme B-specific fragments of DNA-PK_{CS} and NuMA are generated in Fas-negative target cells attacked by lymphokine-activated killer (LAK) cells.

To determine whether granzyme B-specific fragments were generated during lymphocyte-induced cytotoxicity, we used the Fas-negative cell line K562 as targets for LAK cells (McGahon *et al.* 1995; Topalian *et al.*, 1989). To permit biochemical analysis of cleaved proteins in target cells, effector:target cell ratios of 5:1 were used. The signature 89kDa caspase-3 fragment of PARP was generated during LAK-induced target cell death (FIG. 7, lane 3). PARP cleavage was entirely abolished by 100μM Ac-DEVD-CHO (FIG. 7, lane 4). This was consistent with results observed when target cell death was initiated with YT cell granule contents (FIG. 6, lane 4). Using anti-C-terminal antibodies, generation of both the 100kDa granzyme B-specific fragment of DNA-

PK_{CS} and the 150-kDa caspase-3-specific fragment were detected during LAK-induced target cell death (FIG. 7, lane 3). In addition, a 120kDa fragment was observed in these cells, consistent with the caspase-3-mediated cleavage of DNA-PK_{CS} at DWVD-G previously observed both *in vitro* and in intact cells (Casciola-Rosen *et al.*, 1996; Song *et al.*, 1996b). While 100μM Ac-DEVD-CHO inhibited the generation of caspase-3-specific fragments of DNA-PK_{CS} by >90%, the 100kDa granzyme B-specific fragment was insensitive to this inhibitor (FIG. 7, lane 4). This data is consistent with that obtained after initiating target cell death with YT cell granule contents (FIG. 6, lane 4 and data not shown). Similar observations were made for the granzyme B-specific NuMA fragment. Together, these data confirm that the novel fragments of DNA-PK_{CS} and NuMA defined in these studies are indeed generated when intact lymphocytes induce target cell cytotoxicity using the granule pathway.

EXAMPLE 9

Cytotoxic lymphocyte granules induce nuclear morphologic changes: Effects of caspase inhibitors.

Several of the downstream substrates for the caspases are directly and efficiently cleaved by granzyme B during lymphocyte granule-induced cytotoxicity even in the presence of caspase inhibitors. It was determined whether these granules also induced morphologic changes in the target cell. HeLa cells were pre-incubated with YT cell granule contents in the presence or absence of 100μM Ac-DEVD-CHO, prior to addition of Ca²⁺, and further incubation. YT cell granule contents induced the rapid onset (<60 min) of prominent surface blebbing (FIG. 8A), followed by nuclear condensation and fragmentation into membrane-bound apoptotic bodies (FIG. 8B). As described for UVB-induced apoptosis (Casciola-Rosen *et al.* 1994), the autoantigens targeted in systemic autoimmune diseases are rapidly redistributed in target cells exposed to YT cell granule contents, such that they become

clustered around the rim of the condensing apoptotic nucleus (FIG. 8A), and then ultimately around apoptotic bodies (FIG. 8B). Granule content-induced surface blebbing, nuclear fragmentation, formation of apoptotic bodies, and characteristic redistribution of nuclear autoantigens was prevented by Ac-DEVD-CHO (Compare FIG. 8B, 8C). However, a prominent diminution in the size of the nucleus (which was accompanied by condensation of chromatin) was induced by granule contents in Ac-DEVD-CHO-treated cells (FIG. 8C); these nuclear changes were not observed when cells were incubated with Ac-DEVD-CHO alone.

EXAMPLE 10

Antibodies Against Autoantigens and Autoantigenic Fragments.

The present invention also relates to polyclonal and monoclonal antibodies raised in response to the autoantigenic fragments disclosed herein. An antibody is specific for an epitope of an autoantigenic fragment if one of skill in the art can use standard techniques to determine conditions under which one can detect an autoantigenic fragment in a Western Blot of a sample from cells of a tissue. The blot can be of a native or denaturing gel as appropriate for the epitope. An antibody is highly specific for an autoantigenic fragment epitope if no nonspecific background binding is visually detectable. An antibody can also be considered highly specific for an autoantigenic fragment if the binding of the antibody can not be competed by random peptides, polypeptides or proteins, but can be competed by the particular autoantigenic fragment, autoantigen, or peptides or polypeptides derived therefrom.

Autoantigenic fragments can be separated from other cellular proteins by use of an immunoaffinity column made with monoclonal or polyclonal antibodies specific for the autoantigen. Additionally, polyclonal or monoclonal antibodies can be raised against a synthetic peptide (usually from about 9 to about 25 amino acids in length) from a portion of an autoantigen or autoantigenic fragment.

Monospecific antibodies are purified from mammalian antisera containing antibodies reactive against the autoantigenic fragment or are prepared as monoclonal antibodies using the technique of Kohler and Milstein (1975, *Nature* 256: 495-497). Monospecific antibody as used

5 herein is defined as a single antibody species or multiple antibody species with homogenous binding characteristics for the autoantigenic fragment. Homogenous binding as used herein refers to the ability of the antibody species to bind to a specific antigen or epitope, such as those associated with the autoantigenic fragment, as described herein.

10 Autoantigenic fragment -specific antibodies are raised by immunizing animals such as mice, rats, guinea pigs, rabbits, goats, horses and the like, with an appropriate concentration of autoantigenic fragment or a synthetic peptide generated from a portion of the autoantigenic fragment with or without an immune adjuvant.

15 Preimmune serum is collected prior to the first immunization. Each animal receives between about 0.1 mg and about 1000 mg of autoantigenic fragment associated with an acceptable immune adjuvant. Such acceptable adjuvants include, but are not limited to, Freund's complete, Freund's incomplete, alum-precipitate, 20 water in oil emulsion containing *Corynebacterium parvum* and RNA. The initial immunization consists of injecting autoantigenic fragment or peptide fragment thereof, preferably in Freund's complete adjuvant, at multiple sites either subcutaneously (SC), intraperitoneally (IP) or both. Each animal is bled at regular intervals, preferably weekly, to 25 determine antibody titer. The animals may or may not receive booster injections following the initial immunization. Those animals receiving booster injections are generally given an equal amount of autoantigenic fragment in Freund's incomplete adjuvant by the same route. Booster injections are given at about three week intervals until maximal titers 30 are obtained. At about 7 days after each booster immunization or about weekly after a single immunization, the animals are bled, the serum collected, and aliquots are stored at about -20°C.

Monoclonal antibodies (mAb) reactive with the autoantigenic fragment are prepared by immunizing inbred mice, 35 preferably Balb/c, with the autoantigenic fragment. The mice are

immunized by the IP or SC route with about 1 mg to about 100 mg, preferably about 10 mg, of the autoantigenic fragment in about 0.5 ml buffer or saline incorporated in an equal volume of an acceptable adjuvant, as discussed herein. Freund's complete adjuvant is preferred. The mice receive an initial immunization on day 0 and are rested for about 3 to about 30 weeks. Immunized mice are given one or more booster immunizations of about 1 to about 100 mg of the autoantigenic fragment in a buffer solution such as phosphate buffered saline by the intravenous (IV) route. Lymphocytes, from antibody positive mice, preferably splenic lymphocytes, are obtained by removing spleens from immunized mice by standard procedures known in the art. Hybridoma cells are produced by mixing the splenic lymphocytes with an appropriate fusion partner, preferably myeloma cells, under conditions which will allow the formation of stable hybridomas. Fusion partners can include, but are not limited to: mouse myelomas P3/NS1/Ag 4-1; MPC-11; S-194 and Sp 2/0, with Sp 2/0 being preferred. The antibody producing cells and myeloma cells are fused in polyethylene glycol, about 1000 mol. wt., at concentrations from about 30% to about 50%. Fused hybridoma cells are selected by growth in hypoxanthine, thymidine and aminopterin supplemented Dulbecco's Modified Eagles Medium (DMEM) by procedures known in the art. Supernatant fluids are collected from growth positive wells on about days 14, 18, and 21 and are screened for antibody production by an immunoassay such as solid phase immunoradioassay (SPIRA) using the autoantigenic fragment as the antigen. The culture fluids are also tested in the Ouchterlony precipitation assay to determine the isotype of the mAb. Hybridoma cells from antibody positive wells are cloned by a technique such as the soft agar technique of MacPherson, 1973, Soft Agar Techniques, in *Tissue Culture Methods and Applications*, Kruse and Paterson, Eds., Academic Press.

Monoclonal antibodies are produced *in vivo* by injection of pristine primed Balb/c mice, approximately 0.5 ml per mouse, with about 2×10^5 to about 6×10^6 hybridoma cells about 4 days after priming. Ascites fluid is collected at approximately 8-12 days after cell transfer

and the monoclonal antibodies are purified by techniques known in the art.

In vitro production of anti-autoantigenic fragment mAb is carried out by growing the hybridoma in DMEM containing about 2% fetal calf serum to obtain sufficient quantities of the specific mAb. The mAb are purified by techniques known in the art.

Antibody titers of ascites or hybridoma culture fluids are determined by various serological or immunological assays which include, but are not limited to, precipitation, passive agglutination, enzyme-linked immunosorbent antibody (ELISA) technique and radioimmunoassay (RIA) techniques. Similar assays are used to detect the presence of the autoantigenic fragment in body fluids or tissue and cell extracts.

It is readily apparent to those skilled in the art that the herein described methods for producing monospecific antibodies can be utilized to produce antibodies specific for autoantigenic fragment peptide fragments, or full-length autoantigen.

Antibody affinity columns are made, for example, by adding the antibodies to Affigel-10 (Biorad), a gel support which is pre-activated with N-hydroxysuccinimide esters such that the antibodies form covalent linkages with the agarose gel bead support. The antibodies are then coupled to the gel via amide bonds with the spacer arm. The remaining activated esters are then quenched with 1M ethanolamine HCl (pH 8). The column is washed with water followed by 0.23 M glycine HCl (pH 2.6) to remove any non-conjugated antibody or extraneous protein. The column is then equilibrated in phosphate buffered saline (pH 7.3) and the cell culture supernatants or cell extracts containing the autoantigenic fragment are slowly passed through the column. The column is then washed with phosphate buffered saline until the optical density (A_{280}) falls to background, then the protein is eluted with 0.23 M glycine-HCl (pH 2.6). The purified autoantigenic fragment is then dialyzed against phosphate buffered saline.

Levels of an autoantigenic fragment in cells and tissues is quantified by a variety of techniques including, but not limited to, immunoaffinity and/or ligand affinity techniques. Autoantigenic

fragment affinity beads or autoantigenic fragment-specific antibodies are used to isolate ³⁵S-methionine labeled or unlabelled autoantigenic fragment. Labeled autoantigenic fragment is analyzed by SDS-PAGE. Unlabelled autoantigenic fragment is detected by Western blotting,

5 ELISA or RIA assays employing either autoantigenic fragment specific antibodies and/or antiphosphotyrosine antibodies.

Preferred antibodies that bind to an autoantigenic fragment but do not bind to the intact autoantigen or other fragments thereof.

10 Examples of preferred antibodies are those that recognizes a cryptic epitope revealed in the autoantigenic fragment, or an antibody that recognizes a terminal epitope present only in the autoantigenic fragment.

EXAMPLE 11

15 Assay for the Detection of an Autoimmune Condition.

The autoantigenic fragments produced and identified following the teaching of the present invention can be used in a assay to detect the presence of an autoimmune condition. The condition can be the generation of autoantigenic fragments before a disease state evolves,

20 the presence of an autoimmune disease or the lessening of a disease.

The assay is performed on a sample derived from a patient. Most commonly, the sample will be a tissue sample. The presence of autoantigenic fragments can be detected *in situ* or can be partially purified before conducting the assay.

25 To perform an assay within this invention one prepares an autoantigenic fragment. For example, one can prepare an autoantigenic fragment of DNA PK_{CS} by cleaving the protein with granzyme B. The autoantigenic fragment is then used to prepare a monoclonal or polyclonal antibody using any of the methods widely

30 known and used in the art.

The antibody can then be used to qualify or quantify the amount of autoantigenic fragment present in the sample. This can be done by numerous techniques known in the art including using antibody

detectably labeled with ^{125}I , gold, enzyme or other known labels.

Alternatively, a detectable label can be carried on a second antibody specific for the first. The amount of autoantigenic fragment found is quantitatively or qualitatively compared to the amount of found on

5 control cells. A change in the former relative to the latter is indicative of whether an autoimmune disease state is present, is progressing or is reduced.

In an alternative form of the assay one can treat cells as described herein and then isolate the autoantigenic fragments present
10 in treated and control cells. The preparations can be made as crude cell extracts, membrane or intracellular fractions of the cells or after purification steps, *e.g.*, chromatography, precipitation or affinity isolation steps. Crude, partially or highly purified preparations can be analyzed for autoantigenic fragment content, *e.g.*, by using antibodies
15 specific for the autoantigenic fragment.

In another form of the assay, an autoantigenic fragment is used to determine the presence or absence of an autoantibody in a patient as an indication of the presence or absence of an autoimmune condition. The use of particular types or autoantigenic fragments can also indicate
20 the type of autoimmune condition. The autoantibody to be assayed for can be present in the serum or a tissue sample of the patient. An autoantibody can be detected *in situ* or after some purification of immunoglobins from the patient. In one format of the assay, the autoantigenic fragment can be fixed to a support, an autoantibody
25 present in a sample is then contacted with the fragment to permit binding of the autoantibody to the autoantigenic fragment. After appropriate washing, the presence of bound autoantibody can be detected by methods available in the art, including the use of a labeled second antibody against the antibodies from the patient.

30 In any assay it can be advantageous to devise an internal control so that the results of different runs of assays can be compared to each other. A cellular protein that is unrelated to the autoantigenic fragment and present in relatively constant amounts in the cells used in the assay can serve as an internal control.

The assays described above are exemplary of all of the assays within the scope of the present invention. Those of skill in the art can use the autoantigenic fragments and antibodies of this invention in many assay formats known or developed in the art.

5

EXAMPLE 12

Tolerizing a Patient to the Presence of an Autoantigenic fragment.

The present invention provides a method of tolerizing a patient to the future *in vivo* generation of compounds that are normally autoantigenic. This method can be prophylactic.

A patient diagnosed to be at risk of developing an autoimmune response is identified. A sample of the tissue to which the autoimmune response is possible is isolated from the patient. Autoantigenic fragments that can be generated from the tissue are then identified. The autoantigenic fragments are administered to the patient in pharmaceutically acceptable carriers without an adjuvant to induce low-zone tolerance.

Tolerization typically involves purification of relevant autoantigenic fragments in a non-aggregated form. In particular embodiments, autoantigenic fragments of DNA Pk_{CS}, NuMA or PARP are generated by the action of granzyme B.

The autoantigenic fragments can also be present in a mixture. One such mixture can be the product of the application of the contents of granules to a sample of tissue to which a potential autoimmune response is diagnosed. In that case, the autoantigenic fragments are produced in the mixture by the action of the granule contents, including granzyme B.

In any case, the autoantigenic fragments are administered at a low dose as chosen by a skilled physician or veterinarian to induce a low-zone tolerance in the patient. Once tolerization of the patient is achieved, if the normally autoantigenic fragments are produced in the tissue *in vivo*, the immune system will not mount a response against

them and the occurrence of an autoimmune disease state can be avoided or the severity reduced.

EXAMPLE 13

5 Treatment for Malignant Cells.

The present invention also provides a method of generating an autoimmune response against certain cells in a patient. For example, one can induce an autoimmune response against malignant cells in a patient that would benefit from such a response.

- 10 For example, one can isolate a sample of malignant cells from a patient and contact the cells with the contents of granules or granzyme B. The action of the granule contents or granzyme B on the cells can produce autoantigenic fragments from autoantigens present in the cells. The resulting mixture can then be administered to the patient.
- 15 In this case, it is preferred that an adjuvant be administered with the autoantigenic fragments.

- Those components of the mixture that are not altered to produce a new antigen will be recognized by the immune system as self molecules. However, those components that are altered to produce new
- 20 autoantigenic fragments will be seen by the immune system as non-self and an immune response will be generated against them.

- Because the autoantigenic fragments produced in this way are the same as those produced *in vivo* by the action of CTLs, NK and LAK cells on the malignant cells, the immune system is stimulated to
- 25 generate a response against the malignant cells. Therefore, the present invention provides a method to heighten or stimulate the natural immune system processes to act against particular types of cells such as malignant cells. The method is particularly advantageous because the *in vivo* production of the autoantigenic fragments from, *e.g.*, malignant
- 30 cells, can occur at rates too low to stimulate the immune system, or at rates that can lead to a tolerization of the immune system.

EXAMPLE 14

Pharmaceutical Compositions.

Pharmaceutically useful compositions comprising autoantigenic fragments of the present invention can be formulated according to known methods such as by the admixture of a pharmaceutically acceptable carrier. Examples of such carriers and methods of formulation can be found in Remington's Pharmaceutical Sciences. To form a pharmaceutically acceptable composition suitable for effective administration, such compositions will contain an effective amount of the inhibitor.

Therapeutic, prophylactic or diagnostic compositions of the invention are administered to an individual in amounts sufficient to treat or diagnose disorders. The effective amount can vary according to a variety of factors such as the individual's condition, weight, sex and age. Other factors include the mode of administration. The appropriate amount can be determined by a skilled physician

The pharmaceutical compositions can be provided to the individual by a variety of routes such as subcutaneous, topical, oral and intramuscular.

The term "chemical derivative" describes a molecule that contains additional chemical moieties which are not normally a part of the base molecule. Such moieties can improve the solubility, half-life, absorption, etc. of the base molecule. Alternatively the moieties can attenuate undesirable side effects of the base molecule or decrease the toxicity of the base molecule. Examples of such moieties are described in a variety of texts, such as Remington's Pharmaceutical Sciences.

Compositions including autoantigenic fragments identified according to the methods disclosed herein can be used alone at appropriate dosages. Alternatively, co-administration or sequential administration of other agents can be desirable.

The present invention also provides a means to obtain suitable topical, oral, systemic and parenteral pharmaceutical formulations for use in the methods of treatment of the present

invention. The compositions containing autoantigenic fragments identified according to this invention as the active ingredient can be administered in a wide variety of therapeutic dosage forms in conventional vehicles for administration. For example, the

5 compositions can be administered in such oral dosage forms as tablets, capsules (each including timed release and sustained release formulations), pills, powders, granules, elixirs, tinctures, solutions, suspensions, syrups and emulsions, or by injection, as appropriate. Likewise, they can also be administered in intravenous (both bolus and

10 infusion), intraperitoneal, subcutaneous, topical with or without occlusion, or intramuscular form, all using forms well known to those of ordinary skill in the pharmaceutical arts.

Advantageously, autoantigenic fragments of the present invention can be administered in a single daily dose, or the total daily

15 dosage can be administered in divided doses of two, three or four times daily.

Furthermore, compounds for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal routes, using those forms of transdermal

20 skin patches well known to those of ordinary skill in that art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

For combination treatment with more than one active

25 agent, where the active agents are in separate dosage formulations, the active agents can be administered concurrently, or they each can be administered at separately staggered times.

The dosage regimen utilizing compositions of the present invention is selected in accordance with a variety of factors including

30 type, species, age, weight, sex and medical condition of the patient; whether the treatment is prophylactic or therapeutic; the severity of the condition to be treated; the route of administration; the renal, hepatic and cardiovascular function of the patient; and the particular compound thereof employed. A physician or veterinarian of ordinary skill can

35 readily determine and prescribe the effective amount of the drug

- required to prevent, counter or arrest the progress of the condition. Optimal precision in achieving concentrations of compounds of this invention, including purified autoantigenic fragments, within the range that yields efficacy without toxicity requires a regimen based on the
- 5 kinetics of the compound's availability to target sites. This involves a consideration of the distribution, equilibrium, and elimination of a compound.

EXAMPLE 15

10 Autoantigens Cleaved by Granzyme B

- Granzyme B efficiently cleaves three caspase-3 substrates generating unique fragments not generated during any other form of cell death. To determine whether the generation of unique autoantigen fragments by granzyme B was a universal feature of autoantigens, a
- 15 wide range of autoantigens were tested for cleavage by granzyme B *in vitro* and *in vivo*. It was determined that despite their diverse structure, distribution and function, >70% of the autoantigens described in systemic autoimmune diseases are efficiently cleaved by granzyme B and unique fragments are produced. In contrast, granzyme B does not
- 20 generate unique fragments in all the non-autoantigen molecules tested. A panel of autoantigens discovered to be susceptible to cleavage by granzyme B, are listed in Table 3 along with the sites of cleavage.

- The granzyme B cleavage sites in autoantigens were defined. In all cases, the tetrapeptide sequence immediately adjacent to
- 25 the cleavage site was highly conserved. The susceptibility to granzyme B cleavage is therefore a specific, unifying feature of these otherwise unrelated molecules. Furthermore, the ability of granzyme B to generate unique fragments of these antigens indicates that granzyme B plays a mechanistic role in selectively producing the fragments of these
- 30 molecules against which autoimmune responses are initiated. These results highlight a potential role for the cytotoxic lymphocyte granule-induced death pathway in initiation and propagation of autoimmunity.

To determine whether susceptibility to direct cleavage by granzyme B was an isolated feature of PARP, NuMA and DNA-PK_{cs} shown above, or whether it was a more general feature of autoantigens, a variety of well-defined autoantigens selected from across the spectrum of systemic autoimmune diseases were tested for susceptibility to cleavage by purified granzyme B. The efficiency of cleavage of the substrates by purified granzyme B was also noted (Table 3). Initially, a series of autoantibodies of well-defined specificity were used to immunoblot lysates of HeLa cells that had been incubated *in vitro* with or without granzyme B. Lysates were pre-treated with iodoacetamide (IAA) to prevent interference by endogenous caspase activity.

Interestingly, several autoantigens that have previously been shown to be cleaved by caspases during apoptosis were also efficiently cleaved by granzyme B. These substrates included U1-70kDa, topoisomerase-1, SRP-72, PARP and NOR-90. In each case, unique fragments were generated. Mi-2, PMS2 and Ki-67 were also identified as additional autoantigens that are cleaved into distinct, different sets of fragments both by caspases and by granzyme B in these lysates.

Efficient cleavage and generation of novel fragments was also confirmed using purified granzyme B to cleave *in vitro* translated substrates in the case of U1-70kDa, PARP, topoisomerase-1, PMS2, and Mi-2. The cleavage efficiency (k_{cat}/K_m) of those substrates by granzyme B was determined using defined amounts of purified protease to cleave either endogenous substrates in cell lysates or radiolabeled substrates expressed by *in vitro* transcription/translation, or both. Previous studies have demonstrated that equivalent results are obtained using either form of substrate (Andrade *et al.*, 1998). k_{cat}/K_m values varied between $1.39 \times 10^4 \text{ M}^{-1}\text{s}^{-1}$ (PMS2) and $1.6 \times 10^6 \text{ M}^{-1}\text{s}^{-1}$ (topoisomerase-1) (see Table 3).

Previous studies have identified several autoantigens that are not susceptible to cleavage by caspases during apoptosis (Casciola-Rosen *et al.*, 1995 & 1996). Many of these autoantigens were efficiently cleaved by granzyme B. These molecules included fibrillarin, PMS1, CENP-B, Ku-70, La and RNA polymerase II large subunit. The efficiency of cleavage of these substrates by granzyme B varied between $5.9 \times 10^3 \text{ M}^{-1}\text{s}^{-1}$

(CENP-B) and $8 \times 10^4 \text{ M}^{-1}\text{s}^{-1}$ (La) (See Table 3). Interestingly, several ribonucleoprotein autoantigens were not susceptible to cleavage by either granzyme B or caspases. These included Ro52kDa and 60kDa, ribosomal protein P, histones and Sm proteins.

5 Susceptibility to cleavage by granzyme B was a highly specific feature of autoantigens. None of 18 different human non-autoantigens tested was cleaved by granzyme B. The precursors of caspases 3 and 7 are not known to be autoantigens, but are efficiently cleaved by granzyme B. Interestingly, these substrates are cleaved at the same sites by
10 granzyme B and caspase-8, generating identical fragments (see below).

 Since some well-defined autoantibodies do not recognize their antigens by immunoblotting, the susceptibility of radiolabeled endogenous substrates to cleavage by granzyme B *in vitro* was assessed. To perform these studies, HeLa cells were radiolabeled with [^{35}S]
15 methionine/cysteine, and proteins were immunoprecipitated using human autoantibodies. Protein A-agarose beads containing washed precipitated proteins were resuspended in buffer supporting the activity of granzyme B, and incubated in the absence or presence of added purified granzyme B. Reaction products were visualized using SDS-
20 PAGE and fluorography. To confirm the validity of this approach, several different autoantigens known to be cleaved by granzyme B, as well as several autoantigens and non-autoantigens that are not cleaved by granzyme B, were tested. Using autoantibodies that both immunoblot and immunoprecipitate, the cleavage profiles obtained using granzyme B
25 to cleave molecules in lysates (followed by detection with immunoblotting), and after immunoprecipitation (from [^{35}S]methionine-labeled HeLa lysate) were compared. Identical results were obtained using these 2 methods for cleaved autoantigens (topoisomerase-1, Mi-2, RNA polymerase II large subunit, Ku-70, PARP, and NOR-90), uncleaved autoantigens (Ku-80, Ro 60k), and control
30 substrates (β -tubulin, vinculin). Using this immunoprecipitation approach, it was demonstrated that several additional autoantigens (PMScl, RNA polymerase I large subunit, histidyl tRNA synthetase, isoleucyl tRNA synthetase and alanyl tRNA synthetase) were cleaved by

granzyme B, generating unique fragments (Table 3). Of note, all these additional autoantigens (with the exception of RNA polymerase I) are targeted in autoimmune myositis. However, two other tRNA synthetases (threonyl tRNA synthetase and glycyl tRNA synthetase) that are also autoantigens in autoimmune myositis, were not cleaved using this approach.

Thus, in addition to the three autoantigens described above to be cleaved by both caspase-3 and granzyme B, these results identify an additional 7 autoantigens have been identified that are cleaved by both proteases but at different sites. Furthermore, another 10 autoantigens are cleaved exclusively by granzyme B, and not by caspases. Therefore, 20 autoantigens targeted across the spectrum of human systemic autoimmune diseases are efficiently cleaved by granzyme B, generating unique fragments not observed during other forms of cell death. (Table 3).

To confirm that similar autoantigen fragments are generated in intact cells during granule-induced cell death, K562 cells were exposed to YT cell granule contents in the presence of Ca^{2+} , and the biochemical status of the autoantigens were analyzed by immunoblotting. In those cases where autoantigens are substrates for both caspases and granzyme B, both fragments were generated (U1-70kDa, PARP, Mi-2, topoisomerase-1, Ki-67). Autoantigens known to be cleaved only by granzyme B were indeed cleaved in the K562/YT granule system and the granzyme B-specific fragments of Ku-70, PMS-1 and RNA polymerase II large subunit were generated.

It was shown that granzyme B-specific fragments of DNA-PKcs are generated during cytotoxic lymphocyte granule-induced target cell death. A similar approach was used to determine whether other autoantigens cleaved by granzyme B *in vitro* are also cleaved during killing of intact Fas-negative target cells by lymphokine-activated killer (LAK) cells. Granzyme B-specific fragments of Mi-2, U1-70kDa, topoisomerase-1, PMS-1 and SRP-72 as well as Ku-70, RNA polymerase II and Ki-67, are generated during this form of cell death and were identified by immunoblotting with appropriate antibodies. In the cases of Mi-2, U1-70kDa, SRP-72 and topoisomerase-1, which are all

susceptible to direct cleavage by both caspase-3 and granzyme B, the amounts of granzyme B-specific fragments of these antigens appear to be determined by the relative efficiency of cleavage by the two proteases. Thus, granzyme B-specific fragments of Mi-2 and topoisomerase-1 (which are efficiently cleaved by granzyme B) were clearly observed. In contrast, very low levels of granzyme B-specific fragments of U1-70kDa, SRP-72 or the less efficiently cleaved sites on topoisomerase-1 (72 and 75kDa) were observed in the intact cell killing assay unless caspases were inhibited by adding Ac-DEVD-CHO. The cleavage of PMS1, which is a substrate for granzyme B but not for caspases, was unaffected by Ac-DEVD-CHO.

EXAMPLE 16

Specificity of Granzyme B Cleavage of Autoantigens

Granzyme B is a serine protease whose specificity has been defined using a positional scanning combinatorial tetrapeptide library. The protease has a preference for I, V or L in P₄, E, G, S in P₃, and P, S, N, A, Q, H, T, V, E, D, in P₂, with a preference for D in P₁. The sizes of the fragments generated by granzyme B cleavage and the cleavage specificity was used to identify likely cleavage sites. Since granzyme B prefers Asp in the P₁ position, and does not tolerate Ala at that site, site-directed mutagenesis was employed to make a series of Asp-Ala substitutions in several of the granzyme B substrates. The effects of each mutation on the efficiency of cleavage by the protease was assessed.

The granzyme B cleavage sites in PARP and DNA-PKcs have been defined. Using the above approach, the granzyme B cleavage sites in fibrillarin, Mi-2, topoisomerase 1, PMS1, PMS2, and U1-70kDa were also defined (Table 3). Interestingly, 10 of 11 of these cleavage sites contain P (7), A (2), or S (1) in the P₂ position, which are preferred by granzyme B but are poorly tolerated by group III caspases. Furthermore, 4 cleavage sites also contain G or S in P₃. These residues are also not tolerated by group III caspases. Using fragment sizes to predict likely granzyme B cleavage sites in other autoantigens, likely

cleavage sites were also identified in these proteins. In every case, these cleavage sites contained residues in P2 and/or P3 which are preferred by granzyme B, but are not tolerated by group III caspases (Table 3).

To confirm that these substrates were not cleaved by group
 5 III caspases, the substrates were incubated with 50nM purified caspase-
 8. Cleavage assays were performed in HeLa cell lysates in which
 endogenous caspases had first been irreversibly inactivated by 1mM
 iodoacetamide (4°C for 15 min), prior to addition of 5mM DTT to facilitate
 exogenous caspase-8 activity. No cleavage of topoisomerase-1, Mi-2, U1-
 10 70kDa, PARP, Ku-70, RNA polymerase II large subunit, SRP-72, NuMA
 or Ki-67 occurred. Caspase-8 was indeed active in these lysates, as
 evidenced by robust caspase-3 cleavage products of PARP and U1-70kDa
 seen when the IAA 'poisoning' step was omitted. Radiolabeled PMS1
 and PMS2 (generated by IVTT) were also not cleaved by purified caspase-
 15 8.

It was demonstrated that 20 of 28 autoantigens tested were
 susceptible to efficient cleavage by granzyme B, generating unique
 fragments. Although the cleaved molecules differ markedly in
 subcellular location, function, and extended primary sequence, they
 20 share 2 features: (i) all are autoantigens targeted by a high titer
 autoantibody response in human autoimmune diseases, including SLE,
 Sjogren's syndrome, diffuse and limited scleroderma, and autoimmune
 myositis; and (ii) molecules are unified by containing a granzyme B
 cleavage site not susceptible to cleavage by caspase-8 (see below).
 25 Interestingly, autoantibodies against the precursors of caspases 3 and 7
 (which are cleaved by granzyme B and caspase-8 cleave at the same sites
 generating identical fragments) have not been found in >1000
 autoimmune sera screened by immunoblotting.

The status of a molecule as an autoantigen and its unique
 30 susceptibility to cleavage by granzyme B but not by caspase-8, are
 therefore highly related ($P < 0.0001$; Chi-square analysis). The positive
 predictive value of susceptibility to unique cleavage by granzyme B and
 status as an autoantigen is 100% for these 48 substrates, while the
 negative-predictive value is 73%, indicating that additional mechanisms
 35 play a role in selection of some molecules as autoantigens. It is

noteworthy that most of the uncleaved molecules are nucleoprotein complexes (e.g. components of nucleosomes and snRNPs).

The granzyme B cleavage sites in several molecules are highly conserved, even in drosophila and yeast. This striking
 5 conservation of sequence at granzyme B cleavage sites in organisms in which cytotoxic lymphocytes had not yet evolved implies that an important, as yet undefined function is served by these regions. This new, extended family of granzyme B substrates therefore provide a powerful tool with which to explore the evolution and biological functions
 10 of the aspartic acid-specific apoptotic proteases, and to probe the mechanisms of cytotoxic lymphocyte granule-mediated cell death.

Human systemic autoimmune diseases represent a highly complex disease spectrum, with numerous variables affecting individual susceptibility, initiation, and tissue targets. By
 15 demonstrating that the autoantigens targeted across the spectrum of these diseases are unified by their susceptibility to efficient cleavage by granzyme B, with the generation of unique fragments not generated during any other form of cell death, these studies focus attention on the role of the cytotoxic lymphocyte granule pathway in initiation of
 20 autoimmunity. Where substrates are cleaved by both caspases and granzyme B, generation of unique granzyme B fragments is dependent on relative inhibition of the caspases. Without wishing to be bound by a particular theory, it is therefore proposed that during pro-immune intracellular infections occurring in a microenvironment in which
 25 caspase activity is under relative inhibition, production of unique granzyme B fragments is favored. In susceptible individuals in whom clearance of apoptotic material is impaired, suprathreshold amounts of these fragments accumulate and are effectively captured and presented by dendritic cells. The resulting immune response is directed against
 30 products of CTL granule-induced death, generating an autoamplifying injury characteristic of these self-sustaining diseases.

EXAMPLE 17

Screening for Candidate Agents for Treatment

The assays described herein can be adapted for screening for candidate agents for the prophylactic or therapeutic treatment of autoimmune disease, cancer, or the symptoms of such diseases. In an exemplary format, a candidate agent is contacted with both an uncleaved autoantigen and the contents of a lymphocyte granule, a granule enzyme, or granzyme B. The granule enzyme or granzyme B can be prepared in varying degrees of purity. The autoantigen should be a substrate cleavable by the granule contents or, if a purified or partially purified enzyme is used, a substrate for the particular enzyme. Once contacted, one can monitor the cleavage of the autoantigen into autoantigenic fragments. If desired, one can run a control assay with no candidate agent, or a known inhibitor of the enzyme, in parallel. The production of autoantigenic fragments can be monitored by a variety of means known in the art including antibody capture of the epitopes produced through cleavage, the loss of epitopes that span the cleavage site, separation of cleavage products through chromatography or electrophoresis and other techniques known and used in the art or developed subsequently in the art of detection. A screening assay can be quantitative or qualitative.

A candidate agent can be a chemical compound, organic or inorganic, or a biochemical compound including proteins, peptides, glyco-proteins or peptides, polysaccharides or other macromolecules.

A candidate agent that decreases the rate or the amount of cleavage of the autoantigen to autoantigenic fragments is referred to as an inhibitor of the process. Candidate agents can be studied to determine their suitability for application in the treatment of animals and humans by methods and procedure recognized in the art of pharmaceutical sciences. Those candidates which through testing are shown to have appropriate efficacy and an acceptable safety profile are used in the prophylactic or therapeutic treatment of patients.

REFERENCES

- Andrade, F., Roy, S., Nicholson, D., Thornberry, N., Rosen, A., and Casciola-Rosen, L. 1998. Granzyme B Directly and Efficiently Cleaves
5 Several Downstream Caspase Substrates: Implications for CTL-Induced Apoptosis *Immunity* 8:451-460.
- Bach, J.F. and S. Koutouzov. (1997). New clues to systemic lupus. *Lancet* 350:11-11.
- 10 Beidler, D.R., Tewari, M., Friesen, P.D., Poirier, G., and Dixit, V.M. (1995). The baculovirus p35 protein inhibits Fas- and tumor necrosis factor-induced apoptosis. *J.Biol.Chem.* 270, 16526-16528.
- Bockenstedt, L.K., R.J. Gee, and M.J. Mamula. (1995). Self-peptides in the initiation of lupus autoimmunity. *J.Immunol.* 154:3516-3524.
- 15 Bump, N.J., Hackett, M., Hugunin, M., Seshagiri, S., Brady, K., Chen, P., Ferenz, C., Franklin, S., Ghayur, T., Li, P., Licari, P., Mankovich, J., Shi, L.F., Greenberg, A.H., Miller, L.K., and Wong, W.W. (1995). Inhibition of ICE family proteases by baculovirus antiapoptotic protein p35. *Science* 269, 1885-1888.
- 20 Burlingame, R.W., R.L. Rubin, R.S. Balderas, and A.N. Theofilopoulos. (1993). Genesis and evolution of antichromatin autoantibodies in murine lupus implicates T-dependent immunization with self-antigen. *J.Clin.Invest.* 91:1687-1696.
- Casciola-Rosen, L.A., Anhalt, G.J., and Rosen, A. (1994a).
25 Autoantigens targeted in systemic lupus erythematosus are clustered in two populations of surface structures on apoptotic keratinocytes. *J. Exp. Med.* 179, 1317-1330.
- Casciola-Rosen, L.A., Miller, D.K., Anhalt, G.J., and Rosen, A. (1994b).
30 Specific cleavage of the 70-kDa protein component of the U1 small nuclear ribonucleoprotein is a characteristic biochemical feature of apoptotic cell death. *J.Biol.Chem.* 269, 30757-30760.

- Casciola-Rosen, L.A., Anhalt, G.J., and Rosen, A. (1995). DNA-dependent protein kinase is one of a subset of autoantigens specifically cleaved early during apoptosis. *J.Exp.Med.* 182, 1625-1634.
- 5 Casciola-Rosen, L.A., Nicholson, D.W., Chong, T., Rowan, K.R., Thornberry, N.A., Miller, D.K., and Rosen, A. (1996). Apopain/CPP32 cleaves proteins that are essential for cellular repair: A fundamental principle of apoptotic death. *J.Exp.Med.* 183, 1957-1964.
- 10 Casciola-Rosen, L. and A. Rosen. (1997). Ultraviolet light-induced keratinocyte apoptosis: A potential mechanism for the induction of skin lesions and autoantibody production in LE. *Lupus* 6:175-180.
- Casiano, C.A., Martin, S.J., Green, D.R., and Tan, E.M. (1996). Selective cleavage of nuclear autoantigens during CD95 (Fas/APO-1)-mediated T cell apoptosis. *J.Exp.Med.* 184, 765-770.
- 15 Chinnaiyan, A.M. and Dixit, V.M. (1996a). The cell-death machine. *Curr.Biol.* 6, 555-562.
- Chinnaiyan, A.M., Hanna, W.L., Orth, K., Duan, H.J., Poirier, G.G., Froelich, C.J., and Dixit, V.M. (1996b). Cytotoxic T-cell-derived granzyme B activates the apoptotic protease ICE-LAP3. *Curr.Biol.* 6, 897-899.
- 20 Darmon, A.J., Ley, T.J., Nicholson, D.W., and Bleackley, R.C. (1996). Cleavage of CPP32 by granzyme B represents a critical role for granzyme B in the induction of target cell DNA fragmentation. *J.Biol.Chem.* 271, 21709-21712.
- 25 Darmon, A.J., Nicholson, D.W., and Bleackley, R.C. (1995). Activation of the apoptotic protease CPP32 by cytotoxic T-cell-derived granzyme B. *Nature* 377, 446-448.
- Deveraux, Q.L., Takahashi, R., Salvesen, G.S. and Reed, J.C. (1997). X-linked IAP is a direct inhibitor of cell-death proteases. *Nature* 38, 300-304.

- Diamond, B., J.B. Katz, E. Paul, C. Aranow, D. Lustgarten, and M.D. Scharff. (1992). The role of somatic mutation in the pathogenic anti-DNA response. *Ann.Rev.Immunol.* 10:731-757.
- 5 Duan, H., Orth, K., Chinnaiyan, A.M., Poirier, G.G., Froelich, C.J., He, W.-W., and Dixit, V.M. (1996). ICE-LAP6, a novel member of the ICE/Ced-3 gene family, is activated by the cytotoxic T cell protease granzyme B. *J.Biol.Chem.* 271, 16720-16724.
- 10 Fernandes-Alnemri, T., Armstrong, R.C., Krebs, J., Srinivasula, S.M., Wang, L., Bullrich, F., Fritz, L.C., Trapani, J.A., Tomaselli, K.J., Litwack, G., and Alnemri, E.S. (1996). *In vitro* activation of CPP32 and Mch3 by Mch4, a novel human apoptotic cysteine protease containing two FADD-like domains. *Proc.Natl.Acad.Sci.USA* 93, 7464-7469.
- 15 Froelich, C.J., Hanna, W.L., Poirier, G.G., Duriez, P.J., D'Amours, D., Salvesen, G.S., Alnemri, E.S., Earnshaw, W.C., and Shah, G.M. (1996a). Granzyme B perforin-mediated apoptosis of jurkat cells results in cleavage of poly(ADP-ribose) polymerase to the 89-kDa apoptotic fragment and less abundant 64-kDa fragment. *Biochem.Biophys.Res.Comm.* 227, 658-665.
- 20 Froelich, C.J., Orth, K., Turbov, J., Seth, P., Gottlieb, R., Babior, B., Shah, G.M., Bleackley, R.C., Dixit, V.M., and Hanna, W. (1996b). New paradigm for lymphocyte granule-mediated cytotoxicity - Target cells bind and internalize granzyme B, but an endosomolytic agent is necessary for cytosolic delivery and subsequent apoptosis. *J.Biol.Chem.* 271, 29073-29079.
- 25 Ghayur, T., Hugunin, M., Talanian, R.V., Ratnofsky, S., Quinlan, C., Emoto, Y., Pandey, P., Datta, R., Huang, Y., Kharbanda, S., Allen, H., Kamen, R., Wong, W., and Kufe, D. (1996). Proteolytic activation of protein kinase C δ by an ICE/CED-3-like protease induces features of apoptosis. *J.Exp.Med.* 184, 2399-2404.

- Greidinger, E.L., Miller, D.K., Yamin, T.-T., Casciola-Rosen, L., and Rosen, A. (1996). Sequential activation of three distinct ICE-like activities in Fas-ligated Jurkat cells. *FEBS Lett.* **390**, 299-303.
- 5 Gu, Y., Sarnecki, C., Fleming, M.A., Lippke, J.A., Bleakley, R.C., and Su, M.S.S. (1996). Processing and Activation of CMH-1 by Granzyme B. *J.Biol.Chem* **271**, 10816-10820.
- Heusel, J.W., Wesselschmidt, R.L., Shresta, S., Russell, J.H., and Ley, T.J. (1994). Cytotoxic lymphocytes require granzyme B for the rapid induction of DNA fragmentation and apoptosis in allogeneic target cells.
- 10 *Cell* **76**, 977-987.
- Irmeler, M., Thome, M., Hahne, M., Schneider, P., Hofmann, B., Steiner, V., Bodmer, J.L., Schröter, M., Burns, K., Mattmann, C., Rimoldi, D., French, L.E., and Tschopp, J. (1997). Inhibition of death receptor signals by cellular FLIP. *Nature* **388**, 190-195.
- 15 Jacobson, M.D., Weil, M., and Raff, M.C. (1997). Programmed cell death in animal development. *Cell* **88**, 347-354.
- Jans, D.A., Jans, P., Briggs, L.J., Sutton, V., and Trapani, J.A. (1996). Nuclear transport of granzyme B (fragmentin 2). Dependence on perforin *in vivo* and cytosolic factors *in vitro* . *J.Biol.Chem.* **271**, 30781-
- 20 30789.
- Krajewska, M., Wang, H.G., Krajewski, S., Zapata, J.M., Shabaik, A., Gascoyne, R., and Reed, J.C. (1997). Immunohistochemical analysis of *in vivo* patterns of expression of CPP32 (Caspase-3), a cell death protease. *Cancer Res.* **57**, 1605-1613.
- 25 Krajewski, S., Gascoyne, R.D., Zapata, J.M., Krajewska, M., Kitada, S., Chhanabhai, M., Horsman, D., Berean, K., Piro, L.D., Fugier-Vivier, I., Liu, Y.J., Wang, H.G., and Reed, J.C. (1997). Immunolocalization of the ICE/Ced-3-family protease, CPP32 (Caspase-3), in non-Hodgkin's lymphomas, chronic lymphocytic leukemias, and reactive lymph nodes.
- 30 *Blood* **89**, 3817-3825.

- Lanzavecchia, A. (1995). How can cryptic epitopes trigger autoimmunity? *J.Exp.Med.* 181,1945-1948.
- Liu, X., Zou, H., Slaughter, C., and Wang, X. (1997). DFF, a heterodimeric protein that functions downstream of caspase-3 to trigger DNA fragmentation during apoptosis. *Cell* 89, 175-184.
- Mamula, M.J. (1993). The inability to process a self-peptide allows autoreactive T cells to escape tolerance. *J.Exp.Med.* 177:567-571.
- Martin, S.J., Amarante-Mendes, G.P., Shi, L.F., Chuang, T.H., Casiano, C.A., O'Brien, G.A., Fitzgerald, P., Tan, E.M., Bokoch, G.M., Greenberg, A.H., and Green, D.R. (1996). The cytotoxic cell protease granzyme B initiates apoptosis in a cell-free system by proteolytic processing and activation of the ICE/CED-3 family protease, CPP32, via a novel two- step mechanism. *EMBO J.* 15, 2407-2416.
- Martin, S.J. and Green, D.R. (1995). Protease activation during apoptosis: Death by a thousand cuts? *Cell* 82, 349-352.
- McGahon, A.J., Nishioka, W.K., Martin, S.J., Mahboubi, A., Cotter, T.G. and Green, D.R. (1995). Regulation of the Fas apoptotic cell death pathway by Abl. *J. Biol. Chem.* 270, 22625-22631.
- Muzio, M., Chinnaiyan, A.M., Kischkel, F.C., O'Rourke, K., Shevchenko, A., Ni, J., Scaffidi, C., Bretz, J.D., Zhang, M., Gentz, R., Mann, M., Krammer, P.H., Peter, M.E., and Dixit, V.M. (1996). FLICE, a novel FADD-homologous ICE/CED-3-like protease, is recruited to the CD95 (Fas/APO-1) death-inducing signaling complex. *Cell* 85, 817-827.
- Nicholson, D.W., Ali, A., Thornberry, N.A., Vaillancourt, J.P., Ding, C.K., Gallant, M., Gareau, Y., Griffin, P.R., Labelle, M., Lazebnik, Y.A., Munday, N.A., Raju, S.M., Smulson, M.E., Yamin, T., Yu, V.L., and Miller, D.K. (1995). Identification and inhibition of the ICE/CED-3 protease necessary for mammalian apoptosis. *Nature* 376, 37-43.

- Nicholson, D.W. and Thornberry, N.A. (1997). Caspases: Killer proteases. *TIBS* 22, 299-306.
- Odake, S., Kam, C.M., Narasimhan, L., Poe, M., Blake, J.T., Krahenbuhl, O., Tschopp, J., and Powers, J.C. (1991). Human and murine cytotoxic T lymphocyte serine proteases: Subsite mapping with peptide thioester substrates and inhibition of enzyme activity and cytolysis by isocoumarins. *Biochemistry* 30, 2217-2227.
- Pinkoski, M.J., Winkler, U., Hudig, D., and Bleackley, R.C. (1996). Binding of granzyme B in the nucleus of target cells. Recognition of an 80 kDa protein. *J.Biol.Chem.* 271, 10225-10229.
- Podack, E. and Konigsberg, P.J. (1984). Cytolytic T cell granules. Isolation, structural, biochemical and functional characterization. *J.Exp.Med.* 160, 695-710.
- Poe, M., Blake, J.T., Boulton, D.A., Gammon, M., Sigal, N.H., Wu, J.K., and Zweerink, H.J. (1991). Human cytotoxic lymphocyte granzyme B: Its purification from granules and the characterization of substrate and inhibitor specificity. *J.Biol.Chem.* 266, 98-103.
- Quan, L.T., Tewari, M., O'Rourke, K., Dixit, V., Snipas, S.J., Poirier, G.G., Ray, C., Pickup, D.J., and Salveson, G.S. (1996). Proteolytic activation of the cell death protease Yama/CPP32 by granzyme B. *Proc.Natl.Acad.Sci.USA.* 93, 1972-1976.
- Radic, M.Z. and M. Weigert. (1994). Genetic and structural evidence for antigen selection of anti-DNA antibodies. *Ann.Rev.Immunol.* 12:487-520.
- Ramage, P., Cheneval, D., Chve, M., Graff, P., Hemmig, R., Heng, R., Kocher, H.P., Mackenzie, A., Memmert, K., Revesz, L., and Wishart, W. (1995). Expression, refolding, and autocatalytic proteolytic processing of the interleukin-1 β -converting enzyme precursor. *J.Biol.Chem.* 270, 9378-9383.

- Salemi, S., A.P. Caporossi, L. Boffa, M.G. Longobardi, and V. Barnaba. (1995). HIVgp120 activates autoreactive CD4-specific T cell responses by unveiling of hidden CD4 peptides during processing. *J.Exp.Med.* 181:2253-2257.
- 5 Sarin, A., Williams, M.S., Alexander-Miller, M.A., Berzofsky, J.A., Zacharchuk, C.M., and Henkart, P.A. (1997). Target cell lysis by CTL granule exocytosis is independent of ICE/Ced-3 family proteases. *Immunity* 6:209-215.
- 10 Sercarz, E.E., P.V. Lehmann, A. Ametani, G. Benichou, A. Miller, and K. Moudgil. (1993). Dominance and crypticity of T cell antigenic determinants. *Ann.Rev.Immunol.* 11:729-766.
- Sercarz, E.E. and S.K. Datta. (1994). Mechanisms of autoimmunization: perspective from the mid-90s. *Curr.Opin.Immunol.* 6:875-881.
- 15 Shi, L.F., Mai, S., Israel, S., Browne, K., Trapani, J.A., and Greenberg, A.H. (1997). Granzyme B (GraB) autonomously crosses the cell membrane and perforin initiates apoptosis and GraB nuclear localization. *J.Exp.Med.* 185, 855-866.
- 20 Shresta, S., MacIvor, D.M., Heusel, J.W., Russell, J.H., and Ley, T.J. (1995). Natural killer and lymphokine-activated killer cells require granzyme B for the rapid induction of apoptosis in susceptible target cells. *Proc.Natl.Acad.Sci.USA* 92, 5679-5683.
- 25 Simitsek, P.D., D.G. Campbell, A. Lanzavecchia, N. Fairweather, and C. Watts. (1995). Modulation of antigen processing by bound antibodies can boost or suppress class II major histocompatibility complex presentation of different T cell determinants. *J.Exp.Med.* 181:1957-1963.
- 30 Song, Q.Z., Burrows, S.R., Smith, G., Lees-Miller, S.P., Kumar, S., Chan, D.W., Trapani, J.A., Alnemri, E., Litwack, G., Lu, H., Moss, D.J., Jackson, S., and Lavin, M.F. (1996a). Interleukin-1b-converting enzyme-like protease cleaves DNA- dependent protein kinase in cytotoxic T cell killing. *J.Exp.Med.* 184, 619-626.

- Song, Q.Z., Lees-Miller, S.P., Kumar, S., Zhang, N., Chan, D.W., Smith, G.C.M., Jackson, S.P., Alnemri, E.S., Litwack, G., Khanna, K.K., and Lavin, M.F. (1996b). DNA-dependent protein kinase catalytic subunit: A target for an ICE-like protease in apoptosis. *EMBO J.* *15*, 3238-3246.
- 5 Srinivasula, S.M., Fernandes-Alnemri, T., Zangrilli, J., Robertson, N., Armstrong, R.C., Wang, L.J., Trapani, J.A., Tomaselli, K.J., Litwack, G., and Alnemri, E.S. (1996). The Ced-3/interleukin 1 β converting enzyme-like homolog Mch6 and the lamin-cleaving enzyme Mch2a are substrates for the apoptotic mediator CPP32. *J.Biol.Chem.* *271*, 27099-10 27106.
- Srinivasula, S.M., Ahmad, M., Otilie, S., Bullrich, F., Banks, S., Wang, Y., Fernandes-Alnemri, T., Croce, C.M., Litwack, G., Tomaselli, K.J., Armstrong, R.C and Alnemri, E.S. (1997). FLAME-1, a novel FADD-like anti-apoptotic molecule that regulates Fas/TNFR1-induced 15 apoptosis. *J.Biol.Chem.* *272*, 18542-18545.
- Talanian, R.V., Yang, X., Turbov, J., Seth, P., Ghayur, T., Casiano, C.A., Orth, K., and Froelich, C.J. (1997). Granule-mediated killing: Pathways for granzyme B-initiated apoptosis. *J.Exp.Med.* *186*, 1323-1331.
- Thome, M., Schneider, P., Hofmann, K., Fickenscher, H., Meinl, E., 20 Neipel, F., Mattmann, C., Burns, K., Bodmer, J.L., Schröter, M., Scaffidi, C., Krammer, P.H., Peter, M.E., and Tschopp, J. (1997). Viral FLICE-inhibitory proteins (FLIPs) prevent apoptosis induced by death receptors. *Nature* *386*, 517-521.
- Thompson, C.B. (1995). Apoptosis in the pathogenesis and treatment of 25 disease. *Science* *267*, 1456-1462.
- Thornberry, N.A. and Molineaux, S.M. (1995). Interleukin-1 β converting enzyme: a novel cysteine protease required for IL-1 β production and implicated in programmed cell death. *Protein Science* *4*, 3-12.

- Thornberry, N.A., Ranon, T.A., Pieterse, E.P., Rasper, D.M., Timkey, T., Garcia-Calvo, M., Houtzager, V.M., Nordstrom, P.A., Roy, S., Vaillancourt, J.P., Chapman, K.T., and Nicholson, D.W. (1997). A combinatorial approach defines specificities of members of the caspase family and granzyme B - Functional, relationships established for key mediators of apoptosis. *J.Biol.Chem* 272, 17907-17911.
- Topalian, S.L., Solomon, D. and Rosenberg, S.A. (1989). Tumor-specific cytolysis by lymphocytes infiltrating human melanomas. *J. Immunol.* 142, 3714-3725.
- 10 Trapani, J.A., Browne, K.A., Smyth, M.J., and Jans, D.A. (1996). Localization of granzyme B in the nucleus - A putative role in the mechanism of cytotoxic lymphocyte-mediated apoptosis. *J.Biol.Chem.* 271, 4127-4133.
- Tschopp, J. (1994). Granzyme B. *Methods Enzymol.* 244, 80-87.
- 15 Wang, S.Y., Miura, M., Jung, Y.K., Zhu, H., Gagliardini, V., Shi, L.F., Greenberg, A.H., and Yuan, J.Y. (1996). Identification and characterization of Ich-3, a member of the interleukin-1 β converting enzyme (ICE)/Ced-3 family and an upstream regulator of ICE. *J.Biol.Chem.* 271, 20580-20587.
- 20 Watts, C. and Lanzavecchia, A. (1993). Suppressive effect of an antibody on processing of T cell epitopes. *J.Exp.Med.* 178:1459-1463.
- White, E. (1996). Life, death, and the pursuit of apoptosis. *Genes Dev.* 10, 1-15.
- Xue, D. and Horvitz, H.R. (1995). Inhibition of the *Caenorhabditis elegans* cell-death protease CED-3 by a CED-3 cleavage site in baculovirus p35 protein. *Nature* 377, 248-251.
- 25 Yamin, T.T., Ayala, J.M., and Miller, D.K. (1996). Activation of the native 45-kDa precursor form of interleukin- 1 β -converting enzyme. *J.Biol.Chem.* 271, 13273-13282.

Young, J.D.-E., Hengartner, H., Podack, E., and Cohn, Z.A. (1986).
Purification and characterization of a cytolytic pore-forming protein
from granules of cloned lymphocytes with natural killer activity. *Cell*
44, 849-859.

5

TABLE I. Comparison of k_{cat}/K_m (catalytic constant) values for the cleavage of different substrates by granzyme B and caspase-3.

The data obtained in time-course experiments were densitometrically scanned and used to calculate the % cleavage of each substrate. These values were fitted to the first order rate equation [% substrate cleaved = $100 \cdot (1 - e^{-((k_{cat} \cdot [E]/K_m) \cdot \text{time})})$] to calculate k_{cat}/K_m . Measurements for each protease-substrate combination were performed at 3 different protease concentrations, enabling experimental variations in k_{cat}/K_m to be assessed.

Substrate	Method of Detection	k_{cat}/K_m ($M^{-1}s^{-1}$)	
		Cleavage by granzyme B	Cleavage by caspase-3
DNA-PK _{cs}	Immunoblotting	$2.5 \pm 0.8 \times 10^6$	$7.5 \pm 0.8 \times 10^6$
NuMA	Fluorography	$5.4 \pm 1.5 \times 10^5$	$5.0 \pm 1.0 \times 10^5$
Caspase-7 ¹	Fluorography	$1.8 \pm 0.6 \times 10^5$	--
Caspase-7 ¹	Immunoblotting	$1.9 \pm 0.6 \times 10^5$	--
Caspase-3	Fluorography	$3.6 \pm 1.0 \times 10^4$	--
Caspase-3	Immunoblotting	$2.3 \pm 0.4 \times 10^4$	--
PARP	Fluorography	$2.3 \pm 1.8 \times 10^4$	$5.0 \pm 0.2 \times 10^6$

15

¹The source of caspase 3 and 7 precursors used in immunoblotting was THP⁻¹ cytosol. Blotting antibodies were specific for either caspase 3 or 7, respectively.

Table II: Different fragments are detected after *in vitro* cleavage of autoantigens with granzyme B versus caspase-3.

The data obtained in FIG. 2, using purified DNA-PK_{CS}, [³⁵S]methionine-labeled PARP, endogenous DNA-PK_{CS} and NuMA, and purified proteases, were used for the tabulation below.

Substrate	Fragments induced after cleavage with		Likely granzyme B cleavage sites
	Granzyme B	Caspase-3	
DNA-PK _{CS} ²	100kDa	150kDa	VDQD3210-G3211
DNA-PK _{CS} ³	250kDa	250kDa	VGPD2698-F2699; DEV2712-N2713
NuMA	175kDa	185kDa	VLGD411-V412
PARP	62kDa	89kDa	VGPD537-S538 (Froelich <i>et al.</i> , 1996a)

² DNA-PK_{CS} fragments were detected by immunoblotting with monoclonal antibody 25-4 or patient sera A.G. and G.A. (which all recognize the C-terminus).

³ DNA-PK_{CS} fragments were detected by monoclonal antibody 18-2 (which recognizes the N-terminus).

202214

TABLE 3: AUTOANTIGENS ARE EFFICIENTLY CLEAVED BY GRANZYME B

	Autoantigen	Cleavage site	$k_{cat}/K_m (M^{-1} \cdot s^{-1})$	Fragments (kDa)
1	DNA-PKcs	VGPD ²⁶⁹⁸ -F	$2.5 \pm 0.8 \times 10^5$	160,100
2	Topoisomerase I	I EAD ¹⁶ -F	$1.6 \pm 0.6 \times 10^5$	97,72
3	NuMA	VATD ¹⁷⁰⁶ -A	$5.4 \pm 1.4 \times 10^5$	175
4	Mi-2	VDPD ¹³¹² -Y	$8.5 \pm 1.9 \times 10^4$	75,72,48
5	La	LEED ²²⁰ -A	$6.1 \pm 1.7 \times 10^4$	21,28
6	PMS1	LTPD ³¹³ -K ISA D ⁴⁹⁶ -E	$6.9 \pm 0.9 \times 10^4$	48
7	Fibrillarin	VGPD ¹⁸⁴ -G	$3.3 \pm 1.9 \times 10^4$	20,17
8	PARP	VDPD ⁵³⁶ -S	$2.3 \pm 1.8 \times 10^4$	72,62,55
9	U1-70kDa	LGND ⁴⁰⁹ -S	$1.3 \pm 0.4 \times 10^4$	60
10	PMS2	VEKD ⁴⁹³ -S	$1.4 \pm 0.6 \times 10^4$	60,45,35
11	Isoleucyl tRNA synthetase (O.J.)	VTPD ⁹⁸² -Q	$7.8 \pm 1.8 \times 10^4$	
12	Histidyl tRNA synthetase (Jo-1)	LGPD ⁴⁸ -E	$2.3 \pm 0.7 \times 10^4$	40
13	Alanyl tRNA synthetase (PL-12)	VAPD ⁶³² -R	1.8×10^4	63,40
14	RNA polymerase I	ICPD ⁴⁴⁸ -M	$1.3 \pm 0.5 \times 10^4$	140
15	KI-67	VCTD ¹⁴⁸¹ -K	$8.1 \pm 2.6 \times 10^3$	168,148
16	PmScl	VEQD ²⁵² -M	$7.5 \pm 1.4 \times 10^3$	85,74,60
17	CENP B	VDSD ⁴⁶⁷ -E	$5.9 \pm 0.2 \times 10^3$	58,40
18	RNA polymerase II	ITPD ³⁷⁰ -P	ND	200
19	SRP 72	VTPD ⁵⁷³ -P	ND	62
20	Ku 70	ISSD ⁷⁸ -R	ND	65

- 21 Tyrosinase ICTD²⁴⁹
 22 E4 VDVD¹²³
 23 Golgi 160
 24 Myosin $\left[\begin{array}{l} \text{VGPD}^{92} \\ \text{SEVD}^{311} \\ \text{I EAD}^{648} \end{array} \right.$

WHAT IS CLAIMED:

1. A composition comprising at least one purified and isolated autoantigenic fragment wherein said fragment is produced from an autoantigen of a pre-apoptotic cell by the action of at least one lymphocyte granule enzyme, wherein said fragment has at least one terminus derived from the cleavage site of said enzyme.
2. The composition of claim 1 wherein the granule enzyme is Granzyme B.
3. The composition of claim 2 wherein the autoantigenic fragment is derived from an autoantigen that is a substrate for a caspase and the fragment is produced by the Granzyme B catalyzed cleavage of said protein at a site that is not cleaved by caspase.
4. The composition of claim 2 comprising an autoantigenic fragment produced by the Granzyme B catalyzed cleavage of an autoantigen selected from the group consisting of DNA PK_{CS}, PARP and NuMA.
5. The composition of claim 4 comprising at least one autoantigenic fragment selected from the group consisting of DNA-PK_{CS} from amino acid 2699 to 4096; DNA-PK_{CS} from amino acid 3211 to 4096; PARP from amino acid 1 to 538; PARP from amino acid 538 to 1004; NuMA from amino acid 412 to 2111 and NuMA from amino acid 1 to 1799.
6. A pharmaceutical composition comprising at least one purified and isolated autoantigenic fragment having at least one terminus derived from a granule enzyme cleavage site, wherein said cleavage site is not cleaved by a caspase, and a pharmaceutically acceptable carrier.
7. The composition of claim 6 wherein the granule enzyme is granzyme B.

8. The composition of claim 6 wherein the autoantigenic fragment is selected from the group of fragments consisting of DNA-PK_{CS} from amino acids 2699 to 4096; DNA-PK_{CS} from amino acids 3211 to 4096; PARP from amino acid 1 to 538; PARP from amino acids 538 to 1004; NuMA from amino acids 412 to 2111 and NuMA from amino acids 1 to 1799.

9. The pharmaceutical composition of claim 6 comprising at least one autoantigenic fragment derived from a malignant cell.

10. A method of treating a patient in need of treatment for an autoimmune disease comprising administering at least one autoantigenic fragment of claim 1.

11. The method of claim 10 wherein the treatment is prophylactic.

12. The method of claim 10 wherein the autoantigenic fragment is selected from the group of fragments consisting of DNA-PK_{CS} from amino acids 2699 to 4096; DNA-PK_{CS} from amino acids 3211 to 4096; PARP from amino acids 538 to 1004; NuMA from amino acids 412 to 2111 and NuMA from amino acids 1 to 1799.

13. The method of claim 10 wherein the method is a method of tolerizing said patient to the presence of said fragment comprising the steps of:

- (a) identifying a target tissue and isolating cells from the tissue,
- (b) providing at least one lymphocyte granule enzyme,
- (c) contacting the cells with said at least one lymphocyte granule enzyme to produce at least one autoantigenic fragment,
- (d) administering said at least one autoantigenic fragment to the patient.

14. The method of claim 13 wherein said at least one lymphocyte granule enzyme provided in step (b) is isolated from the contents of a lymphocyte granule.

5

15. The method of claim 10 for the therapeutic treatment of a patient producing autoantigenic fragments and autoantibodies against the fragments comprising the steps of:

(a) providing an isolated autoantigenic fragment associated with the autoimmune condition in the patient,

(b) contacting the serum of the patient with the autoantigenic fragment under conditions that allow the binding of autoantibodies to said autoantigenic fragment.

16. The method of claim 15 wherein at least a portion of the autoantibodies are removed from the serum of the patient.

17. The method of claim 15 wherein the autoantibodies are bound to the isolated autoantigenic fragment *in vivo*.

18. A method of treating a patient in need of treatment for a malignancy comprising the steps of

(a) providing at least one enzyme of a lymphocyte granule,

(b) isolating malignant cells from the patient,

(c) contacting the malignant cells with the enzyme to produce a mixture containing autoantigenic fragments, and

(d) administering the autoantigenic fragments to the patient.

30

19. An assay for the detection of an autoantigenic fragment in a patient as an indication of the presence or absence of an autoimmune condition in a patient comprising:

(a) providing a sample from the patient,

(b) contacting the sample with an antibody that specifically binds to a cryptic epitope of an autoantigenic fragment, said fragment having at least one terminus derived from a granule enzyme cleavage site,

5 (c) detecting the presence or absence of the binding of the antibody to the autoantigenic fragment as an indication of the presence or absence of an autoimmune condition in a patient.

20. The assay of claim 19 wherein the granule enzyme is
10 granzyme B.

21. An assay for the detection of an antibody that binds an autoantigenic fragment as an indication of the presence or absence of an autoimmune condition in a patient comprising:

15 (a) providing a sample from the patient,

(b) contacting the sample with an autoantigenic fragment having at least one terminus derived from cleavage by a granule enzyme,

(c) detecting the presence or absence of the binding of an
20 antibody in the sample to the autoantigenic fragment as an indication of the presence or absence of an autoimmune condition in the patient.

22. The assay of claim 21 wherein the granule enzyme is
25 granzyme B.

23. A method of making an autoantigenic fragment from an autoantigen comprising the steps of

(a) isolating cells containing at least one autoantigen,
and

30 (b) contacting the cells with a lymphocyte granule enzyme to produce a mixture containing at least one autoantigenic fragment.

24. The method of claim 22 further comprising the step of
35 (c) isolating said at least one autoantigenic fragment.

25. The method of claim 22 wherein step (a) further comprises purifying at least one autoantigen from the cells and step (b) comprises contacting said purified autoantigens with granzyme B.

5

26. The method of claim 25 wherein in step (a) the at least one autoantigen is at least one of DNA-PK_{CS}, PARP and NuMA, and step (b) comprises contacting said at least one autoantigen with granzyme B.

10

27. The method of claim 22 wherein said lymphocyte granule enzyme is isolated from the granules of at least one lymphocyte selected from the group consisting of cytotoxic T lymphocytes (CTL), natural killer cells (NK), lymphokine activated killer cells (LAK) and cells of the YT cell line.

15

28. A method of identifying candidate agents for preventing or treating autoimmune disease symptoms comprising:

a) contacting a test substance with at least one granule enzyme and an autoantigen which is a substrate for said at least one granzyme enzyme;

20

b) monitoring the cleavage of the autoantigen said enzyme into autoantigenic fragments;

c) determining whether the candidate agent alters the production of the autoantigenic fragments;

25

wherein a test substance which inhibits the cleavage is identified as a candidate agent for treating autoimmune diseases.

29. The method of claim 28 wherein the granule enzyme is granzyme B.

30

AUTOANTIGENIC FRAGMENTS, METHODS AND ASSAYS


5

[illegible]

1/17

FIG. 1


Fig 1



2/17

FIG. 2
Fig 2

John R. J. DM-3362
Rosen



3/17

cell " 299660

FIG. 3


DNA	+	-	+	-	+
Granzyme B	-	-	+	+	-
DNA-PK_{cs}	+	+	+	+	-

SP1 — 



Fig 3

4/11
FIG. 4



4
B


5/17

1913	1914	1915	1916	1917	1918	1919	1920	1921	1922	1923	1924	1925	1926	1927	1928	1929	1930	1931	1932	1933	1934	1935	1936	1937	1938	1939	1940	1941	1942	1943	1944	1945	1946	1947	1948	1949	1950	1951	1952	1953	1954	1955	1956	1957	1958	1959	1960	1961	1962	1963	1964	1965	1966	1967	1968	1969	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	2101	2102	2103	2104	2105	2106	2107	2108	2109	2110	2111	2112	2113	2114	2115	2116	2117	2118	2119	2120	2121	2122	2123	2124	2125	2126	2127	2128	2129	2130	2131	2132	2133	2134	2135	2136	2137	2138	2139	2140	2141	2142	2143	2144	2145	2146	2147	2148	2149	2150	2151	2152	2153	2154	2155	2156	2157	2158	2159	2160	2161	2162	2163	2164	2165	2166	2167	2168	2169	2170	2171	2172	2173	2174	2175	2176	2177	2178	2179	2180	2181	2182	2183	2184	2185	2186	2187	2188	2189	2190	2191	2192	2193	2194	2195	2196	2197	2198	2199	2200	2201	2202	2203	2204	2205	2206	2207	2208	2209	2210	2211	2212	2213	2214	2215	2216	2217	2218	2219	2220	2221	2222	2223	2224	2225	2226	2227	2228	2229	2230	2231	2232	2233	2234	2235	2236	2237	2238	2239	2240	2241	2242	2243	2244	2245	2246	2247	2248	2249	2250	2251	2252	2253	2254	2255	2256	2257	2258	2259	2260	2261	2262	2263	2264	2265	2266	2267	2268	2269	2270	2271	2272	2273	2274	2275	2276	2277	2278	2279	2280	2281	2282	2283	2284	2285	2286	2287	2288	2289	2290	2291	2292	2293	2294	2295	2296	2297	2298	2299	2300	2301	2302	2303	2304	2305	2306	2307	2308	2309	2310	2311	2312	2313	2314	2315	2316	2317	2318	2319	2320	2321</
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	--------

File 5

AR copies

Figures



202214

6/17

Fig. 6

Ca ²⁺	+	-	+	+
EDTA	-	+	-	-
Ac-DEVD-CHO	-	-	-	+
Granule Contents	-	+	+	+
	-	-	-	-450

DNA-PKcs






FIG. 7



202214

8/17

FIGURE 8



202214

9/17

FIG. 9A

LOCUS 284337 2101 aa 12-APR-1996
DEFINITION NuMA protein - human.
ACCESSION 284337
PID g284337
DBSOURCE PIR: locus A42184
summary: #length 2101 #molecular-weight 236296 #checksum 8715.
PIR dates: 31-Dec-1993 #sequence_revision 31-Dec-1993 #text_change
12-Apr-1996.
KEYWORDS .
SOURCE human.
ORGANISM Homo sapiens
Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
Vertebrata; Mammalia; Eutheria; Primates; Catarrhini; Hominidae;
Homo.
REFERENCE 1 (residues 1 to 2101)
AUTHORS Compton,D.A., Szilak,I. and Cleveland,D.W.
TITLE Primary structure of NuMA, an intranuclear protein that defines a
novel pathway for segregation of proteins at mitosis
JOURNAL J. Cell Biol. 116 (6), 1395-1408 (1992)
MEDLINE 92176238
REFERENCE 2 (residues 1 to 2101)
AUTHORS Tang,T.K., Tang,C.J., Chen,Y.L. and Wu,C.W.
TITLE Nuclear proteins of the bovine esophageal epithelium. II. The NuMA
gene gives rise to multiple mRNAs and gene products reactive with
monoclonal antibody W1
JOURNAL J. Cell. Sci. 104 (Pt 2), 249-260 (1993)
MEDLINE 93280231
REFERENCE 3 (residues 1 to 2101)
AUTHORS Harborth,J., Weber,K. and Osborn,M.
TITLE Epitope mapping and direct visualization of the parallel,
in-register arrangement of the double-stranded coiled-coil in the
NuMA protein
JOURNAL EMBO J. 14 (11), 2447-2460 (1995)
MEDLINE 95300777
FEATURES Location/Qualifiers
source 1..2101
/organism="Homo sapiens"
/db_xref="taxon:9606"
Protein 1..2101
/product="NuMA protein"

66240" 2993660

10/17

1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	2101	2102	2103	2104	2105	2106	2107	2108	2109	2110	2111	2112	2113	2114	2115	2116	2117	2118	2119	2120	2121	2122	2123	2124	2125	2126	2127	2128	2129	2130	2131	2132	2133	2134	2135	2136	2137	2138	2139	2140	2141	2142	2143	2144	2145	2146	2147	2148	2149	2150	2151	2152	2153	2154	2155	2156	2157	2158	2159	2160	2161	2162	2163	2164	2165	2166	2167	2168	2169	2170	2171	2172	2173	2174	2175	2176	2177	2178	2179	2180	2181	2182	2183	2184	2185	2186	2187	2188	2189	2190	2191	2192	2193	2194	2195	2196	2197	2198	2199	2200	2201	2202	2203	2204	2205	2206	2207	2208	2209	2210	2211	2212	2213	2214	2215	2216	2217	2218	2219	2220	2221	2222	2223	2224	2225	2226	2227	2228	2229	2230	2231	2232	2233	2234	2235	2236	2237	2238	2239	2240	2241	2242	2243	2244	2245	2246	2247	2248	2249	2250	2251	2252	2253	2254	2255	2256	2257	2258	2259	2260	2261	2262	2263	2264	2265	2266	2267	2268	2269	2270	2271	2272	2273	2274	2275	2276	2277	2278	2279	2280	2281	2282	2283	2284	2285	2286	2287	2288	2289	2290	2291	2292	2293	2294	2295	2296	2297	2298	2299	2300	2301	2302	2303	2304	2305	2306	2307	2308	2309	2310	2311	2312	2313	2314	2315	2316	2317	2318	2319	2320	2321	2322	2323	2324	2325	2326	2327	2328	2329	2330	2331	2332	2333	2334	2335	2336	2337	2338	2339	2340	2341	2342	2343	2344	2345	2346	2347	2348	2349	2350	2351	2352	2353	2354	2355	2356	2357	2358	2359	2360	2361	2362	2363	2364	2365	2366	2367	2368	2369	2370	2371	2372	2373	2374	2375	2376	2377	2378	2379	2380	2381	2382	2383	2384	2385	2386	2387</
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	--------

1	mtlhatrgaa	llswvnslhv	adpveavlql	qdcisifikii	drihgteegg	qilkqpvser
61	ldfvcsflqk	nrkhpsspec	lvsaqkvleg	selelakmtm	llyhstms	ksprdweqfe
121	ykiqaelavi	lkfvldhedg	lnlnedlenf	lqkapvpstc	sstfpeelsp	pshgakreir
181	flelqkvass	ssgnnflsgs	paspmgdilg	tpqfgmrrlk	kqladersnr	delelelaen
241	rklltekdaq	iammqgridr	lallnekqaa	splepkelee	lrdknesltm	rihetlkgcq
301	dlkteksqmd	rkinqlseen	gdlsfklref	ashlqqlgda	lnelteehsk	atgewlekqa
361	glekelsaal	qdkkcleekn	eilgqklsq	eehlsqldn	ppqekgevlg	dvlqletlkq
421	eaatlaannt	qlgarvemle	terggqeakl	laerghfeee	kqqlsslitd	lgssisnlsq
481	akeelegasq	ahgarltaqv	asltseletl	natiqqgdqe	laglkqqake	kqaqlaqltq
541	qqeqasqglr	hqveqlsssl	kqkeqqklev	aekgeatrqd	haqqlataae	ereaslrrd
601	aalkqleale	kekaakleil	qqqlgvanea	rdsaqtsvtq	aqrekaelsr	kveelqacve
661	targeqhaeq	aqvaelelql	rseqqkatek	ersaqekdql	qeqlqalkes	lkvtkgslee
721	ekrraadale	eqgrciselk	aetrslveqh	krerkeleee	ragrkglear	llqlgeahqa
781	etevlrrela	eamaaghtae	seceqlvkev	aawrdgyeds	qqeeaqygam	fgeqlmtlike
841	ecekargelq	eakekvagie	shselqisrq	qnklaelhan	laralqqvqe	kevraqklad
901	dlstlqekma	atskevarle	tlvrkageqq	etasrelvke	paragdrqpe	wleeqqgrqf
961	cstqaalgam	ereaeqmgne	lerlraalme	sqqggqeerg	qqerevarlt	qergragadl
1021	alekaarael	emrlqnalne	qrvefatlqe	alahalteke	gkdqelaklr	gleaaqikel
1081	eelrqtvkql	keqlakkeke	hasgsgaqse	aagртеptgp	klealraevs	klegqqcqkqg
1141	eqadslersl	eaerasraer	dsaletlqqg	leekaqelgh	sgsalasaqr	elaaftrtkvq
1201	dhskaedewk	aqvargrgea	erknslissl	eeevsilnrq	vlekegeske	lkrllmaese
1261	ksqkleesca	ccrqrgpatv	pelqnaallc	grrcrasgre	aekgrvasen	lrqeltvsqae
1321	raaelqgskl	awgekkffqk	qalstlgleh	tstqalvsel	lpakhlcqql	qaegaaaeqr
1381	hreeleqskq	aaqglraell	raqrelgeli	plrqlvaeqe	rtaqqlraek	asyaeqlsml
1441	kkahgllaee	nrglgeranl	grqfleveled	qarekyvqel	aavradaetr	laevqreags
1501	tarelevmta	kyegakvkvl	eergfrfeer	qkltaqveel	skkladsdqa	skvqqqklka
1561	vqagqgesqg	eaqrfgaqln	elgaqlsqke	qaaehyklqm	ekakthydak	kqgnqelqeg
1621	lrsleqlqke	nkelraaeer	lghelqqagl	ktkeaeqtr	hltaqvrsle	aqvahadqql
1681	rdlgkfqvat	dalksrepqa	kpqldlsids	ldlsceegtp	lsitsklprt	qpdgtsvpge
1741	paspisqrlp	pkveslesly	ftpiparsqa	plessldslg	dvfldsgrkt	rsarrrrttqi
1801	initmtkkld	veepdsanss	fystrsapas	gaslratsst	qslarlgspd	ygnsallslp
1861	gyrpttrssa	rrsqagvssg	appgrnsfym	gtcqdepeql	ddwnriaelq	qnnrvcpphl
1921	ktcyplesrp	slslgtitde	emktgdpqet	lrrasmqpiq	iaegtgittr	qqrkrvslp
1981	hggpgtpesk	katscfprpm	tprdrhegrk	qstteaakka	apastkqadr	rqsmafsln
2041	tpkklgnsl	rrgaskkals	kaspntrsqt	rrspriattt	asaataaaig	atprakgak
2101	h					

11/17

LOCUS 107227 2115 aa10-NOV-1995
DEFINITION NuMA protein - human.
ACCESSION 107227
PID g107227
DBSOURCE PIR: locus S23647
summary: #length 2115 #molecular-weight 238273 #checksum 4391.
PIR dates: 10-Feb-1994 #sequence_revision 10-Nov-1995 #text_change
10-Nov-1995.
KEYWORDS .
SOURCEhuman.
ORGANISM Homo sapiens
Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
Vertebrata; Mammalia; Eutheria; Primates; Catarrhini; Hominidae;
Homo.
REFERENCE 1 (residues 1 to 2115)
AUTHORS Yang,C.H., Lambie,E.J. and Snyder,M.
TITLE NuMA: an unusually long coiled-coil related protein in the
mammalian nucleus
JOURNAL J. Cell Biol. 116 (6), 1303-1317 (1992)
MEDLINE 92176231
FEATURES Location/Qualifiers
source 1..2115
/organism="Homo sapiens"
/db_xref="taxon:9606"
Protein 1..2115
/product="NuMA protein"

FIG. 10B

1 mtlhatrgaa llswvnslhv adpveavlql qdcsifikii drihgteegq gilkqpvser
 61 ldfvcsflqk nrkhpsspec lvsaqkvleg selelakmtm lllyhstms ksprdweqfe
 121 ykigaelavi lkfvldhedg lnlnedlenf lqkapvpstc sstfpeels pshqakreir
 181 flelqkvass ssgnnflsgs paspmgdilq tpqfqmrrlk kqladersnr delelelaen
 241 rklltekdaq iammqgridr lallnekqaa splepkelee lrdknesltm rlhetlkqcg
 301 dlkteksqmd rkinglseen gdlsfklref ashlqqlqda lnelteehsk atqewlekqa
 361 qlekelsaal qdkkcleekn eilqgklsq eehlsqldn ppqekgevlg dvlqletlkq
 421 eeatlaannt qlgarvemle terggqeakl laerghfeee kqqlsslitd lqssisnlsq
 481 akeeleqasq ahgarltaqv asltselett natigqqdqe laglkqqake kqaqlaqtq
 541 qqeqasqglr hqveqlsssl kqkeqqlkev aekqeatrqd haqqldatae ereaslrer
 601 aalkqleale kekaakleil qqqlqvanea rdsagtsvtg agrekaelsr kveelqacve
 661 tarqeqheaq aqvaelql rseqqkatek ervagekdql geqlqalkes lkvtkgslee
 721 ekrraadale eqgrciselk aetrslveqh krerkeleee ragrkglear lqqlgeahqa
 781 etevlrrela eamaaqhtae seceqlvkev aawreryeds qgeeagygam fqeqlmtlke
 841 ecekargelq eakekvagie shselqisrq qnelaelhan laralqqvqe kevraqklad
 901 dlstlqekma atskevarle tlvrkageeq etasrelvke paragdrqpe wleeqqgrqf
 961 cstqaalgam ereaeqmgne lerlraalme sggqqgeerg qqerevarlt qergraqadl
 1021 alekaarael emrlqnalne qrvefatlqe alahalteke gkdqelaklr gleaaqikel
 1081 eelrqtvkql keqlakkeke hasgsgaqse aagртеptgp klealraevs klegqcqkqg
 1141 eqadslersl eaerasraer dsaletlqq leekaqelgh sqsalasaqr elaafrtkvq
 1201 dhskaedewk aqvargrgea erknsllssl eevevslnrq vlekegeske lkrlvmaese
 1261 ksqkleerlr llqaetasns araaerssal reevqslree aekqrvasen lrgeltsqae
 1321 raeelggelk awqekffqke qalstlqleh tstqalvsel lpakhlcqql qaeqaaekr
 1381 hreelegskq aagglraell raqrelgeli plrqkvaee rtaqqlraek asyaeqlsml
 1441 kkahgllae nrglgeranl grqfleveld garekyvqel aavradaetr laevqreaqs
 1501 tarelevmta kyegakvkvl eergrfgeer qkltaqveql evfqreqtq veelskklad
 1561 sdqaskvqq klkavqaqgg esqqaarlq aqlnelqaql sqkeqaaehy klqmekakth
 1621 ydakkqqnqe lqeqlrsleq lqkenkelra eaerlghelq gaglktkeae qtrhltaqv
 1681 rsleaqvaha dqqlrdlgkf qvatdalksr epqakpql dl sidsldlsce egtplstsk
 1741 lpqrtqpdgts vpgepaspis qrlppkvesl eslyftpipa rsqaplessl dslgdvfdqs
 1801 grktrsarrt ttqiinitmt kkldveepds anssfystrs apasqasla tsstqslarl
 1861 gspdygnsal lslpgyrptt rssarrsqag vsagappgrn sfymgtcde peqlddwnri
 1921 aelqqrnrvc pphlktcyl esrpslsigt itdeemktgd pgetlrrasm qpigiaeagt
 1981 ittrqqrkrv slephqpgt peskkatscf prpmtprdrh egrkqsttea qkkaapastk
 2041 qadrrqsmf silntpkklg nsllrrgask kalskaspnt rsgtrrspri atttasaata
 2101 aaigatprak gkakh

202214

13/17

FIG. 11A

LOCUS 1362789 4096 aa 06-SEP-1996
DEFINITION DNA-activated protein kinase, catalytic subunit - human.
ACCESSION 1362789
PID g1362789
DBSOURCE PIR: locus A57099
summary: #length 4096 #molecular-weight 465420 #checksum 1795.
genetic: #gene GDB:PRKDC ##cross-references GDB:234702
#map_position 8q11.
PIR dates: 27-Oct-1995 #sequence_revision 27-Oct-1995 #text_change
06-Sep-1996.
KEYWORDS DNA binding; DNA recombination; DNA repair; nucleus;
phosphotransferase.
SOURCE human.
ORGANISM Homo sapiens
Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
Vertebrata; Mammalia; Eutheria; Primates; Catarrhini; Hominidae;
Homo.
REFERENCE 1 (residues 1 to 4096)
AUTHORS Sipley,J.D., Menninger,J.C., Hartley,K.O., Ward,D.C., Jackson,S.P.
and Anderson,C.W.
TITLE Gene for the catalytic subunit of the human DNA-activated protein
kinase maps to the site of the XRCC7 gene on chromosome 8
JOURNAL Proc. Natl. Acad. Sci. U.S.A. 92 (16), 7515-7519 (1995)
MEDLINE 95365397
REFERENCE 2 (residues 1 to 4096)
AUTHORS Hartley,K.O., Gell,D., Smith,G.C., Zhang,H., Divecha,N.,
Connelly,M.A., Admon,A., Lees-Miller,S.P., Anderson,C.W. and
Jackson,S.P.
TITLE DNA-dependent protein kinase catalytic subunit: a relative of
phosphatidylinositol 3-kinase and the ataxia telangiectasia gene
product
JOURNAL Cell 82 (5), 849-856 (1995)
MEDLINE 95401275
FEATURES Location/Qualifiers
source 1..4096
/organism="Homo sapiens"
/db_xref="taxon:9606"
Protein 1..4096
/note="DNA-PK-cs"
/product="DNA-activated protein kinase, catalytic subunit"

66240" 29996260

FIG. 11B

1 magsgagvrc slrlrlqetls aadrcgaala ghqlirglgq ecvlsssspav lalqtslvfs
 61 rdfglllvfvr kslnsiefre creeilkflc iflekmggki apysveiknt ctsvytkdra
 121 akckipaldl likllqtfrs srlmdefkig elfskfygel alkkkipdvt lekvyellgl
 181 lgevhpsemi nnaenlfrac lgelktqmts avrepklpvl agclkgllss lcnftksmee
 241 dpqtsreifr fvlkairpqi dlkryavpsa gtrlfalhas qfstclldny vslfeflikw
 301 cahtnvelkk aalsalesfl kgvsnmvakn aemhknklqy emeqfygiir nvdsnnkels
 361 iaargyglfa gpckvinakd vdfmyveliq rckqmfiltqt dtgdyrvyqm psflqsvasv
 421 llyldtvpev ytpvlehlv mquidsfpqys pkmqlvccra ivkvflalaa kgpvlrncis
 481 tvvhqgliri cskpvvlpkg pesesedhra sgevrtdgkwk vptykdyvdl frhlssdqm
 541 mdsiladeaf fsvnsssesl nhlllydefvk svlkivekld ltleiqtvg qengdeapgv
 601 wmiptsdpaa nlhpkpkdf safinlvfc reilpekqae ffepwvysfs yelilqstrl
 661 plisgfykll sitvrnakki kyfegvpsks lkhspedpek yscfalvfkf gkevavkmkq
 721 ykdellascf tfllslphni ieldvrayvp alqmafklgl sytplaevgl naleewsiyi
 781 drhvmqpyyk dilpcldgyl ktsalsdetk nnwevsalsr aaqkgfinkv lkhkktknl
 841 ssneaislee irirvqmlg slggqinknl lvtssdemmm ksyvawdrek rlsfavpfre
 901 mkpviifldvf lprvtelalt asdrqtkvaa cellhsmvmf mlgkatqmp ggggappmyq
 961 lykrtfpvll rladvdvgt rqlieplvmq lihwtfnkk fesqdtvsl eaildgivdp
 1021 vdstlrdfcg rcireflkws ikqitpqgqe kspvntkslf krlyslalhp nafkrlgasl
 1081 afnniyrefr eeelveqfv fealviymes lalahadeks lgtiqqccda idhlcriiek
 1141 khvslnkakk rrlprgfpps aslclldlvk wllahcgrpq tecrhksiel fykfvpllp
 1201 nrspnlwldk vlkeegvsfl intfegggcg qpsgilagpt llylrgpfsl qatlcwldll
 1261 laalecyntf igertvgalg vlgteaqssl lkavaffles iamhdiiaae kcfgtgaagn
 1321 rtspqegery nyskctvvvr imeftttlln tipegwkllk kdlcnthlmr vlvqtlcepa
 1381 sigfnigdvq vmahlpdvcv nlmkalkmsp ykdilethlr ekitagsiee lcavnlygpd
 1441 aqvdrsrllaa vvsackqlhr agllhnlps qstdlhhsdg tellslvykg iapgderqcl
 1501 psldlscqql asgllelafa fggclcerlvs llnpavlst aslgssqgsv ihfshgeyfy
 1561 slfsetinte llknldlav lmqssvdnt kmvsavlnm ldqsfrean qkhqglklat
 1621 tilqhwkcd sawakdsple tkmavllala kilqidssvs fntshgsfpe vfttyislla
 1681 dtkldlhlkg gavltpffft sltggsleel rrvleqliva hfpmqsrefp pgtprfnnyv
 1741 dcmkkfldal elsqspmlle lmtvclreg qhvmeelfqs sfrriarrgs cvtqvglls
 1801 vyemfrkddp rlsftrqsfv drsltlwllh csldalreff stivvdaidv lksrftklne
 1861 stfdtqitkk mgyykildvm ysrlpkddvh akeskingvf hgscitegne ltktlklcy
 1921 daftenmage nqlerrrrly hcaayncas viccvfnelk fyggflfsek peknllifen
 1981 lidlkrrynf pvevevpmer kkkyeirke areaangdsd gpsymsslsy ladstlseem
 2041 sqdfstgvq sysyssqdr patgrfrre qrdptvhddv lelemdelnr hecmaplal
 2101 vkhmhrslgp pqgeedsvpr dlpswmkflh gklgnpivpl nirlflaklv inteevfrpy
 2161 akhwlsplllq laasenngge gihymvveiv atilswtga tptgvpkdev lanrllnflm
 2221 khvfhpkrav frhnleikt lvecwkdcsl ipyrllifekf sgkdpnskd svgiqllgiv
 2281 mandlppydp qcgigsseyf qalvnmsfv rykevyaaaa evlgililryv merknilees
 2341 lcelvakqlk qhantmedkf ivclnkvtks fppladrfmn avfllpkfh gvlktlclev
 2401 vlcrvegmte lyfqlkskdf vqvmhrder qkvclldiik mmpklkpvel relnlpvvef
 2461 vshpsttre qmynilmwih dnyrdpeset dndsgeifkl akdvliqgli denpglqlii
 2521 rnfwshtetrl psntldrlla lnslyspkie vhlslatnf llemtsmspd ypnpmfehpl
 2581 secefqeyti dsdwrfrstv ltpmfvetqa sqgtlqtrtq egslsarwpv agqiratqqq
 2641 hdtltqtad grssfdwltg sstdplvdht spssdsllfa hkrserlqra plksvgpddf
 2701 kkrlglpgde vdnkvkgaag rtdllrlrrr fmrdqekls myarkgvaeg krekeiksel

FIG. 11C

2761 kmkqdaqvvl yrsyrhgdlp diqikhssli tplqavaqrd piiakqlfss lfsgilkemd
2821 kfktilseknn itqkllqdfn rflnttfsff ppfvsciqdi scqhaallsl dpaavsagcl
2881 aslqqpvgir lleallrll paelpakrvr gkarlppdvl rwvelaklyr sigeydvlg
2941 iftseigtkq itqsallaea rsdyseaakq ydealnkqdw vdgepteak dfwelaslde
3001 ynhlaewksl eyctasids enppdlnkiw sepfyqetyl pymirsklkl llqgeadqsl
3061 ltfidkamhg elqkailelh ysqelsllyl lqddvdraky yiqngiqsfm qnyssidvll
3121 hqsrltklqs vqalteiqef isfiskqgnl ssqvplkrll ntwtwnrypda kmddpmniwdd
3181 iitnrcffls kieekltplp ednsmnvdqd gdpsdrmevq egeedissli rsckfsmkknk
3241 midsarkqnn fslamkllke lhkesktrdd wlvswvqsyrc rlshcrsrsq gcseqvltvl
3301 ktvsllldenn vssylxknll afirdqnillg ttyriianal ssepaclaei eedkarrile
3361 lsgsssedse kviaglygra fqlhseavqa aeeeagppsw scgpaagvid aymtladfcd
3421 qqlrkeeena svtdsaclqa ypalvvekml kalklnsnea rlfkprllqi ierypeetls
3481 lmtkeissvp cwqfiswish mvalldkdqa vavqhsveei tdnypqaiyv pfiissesys
3541 fkdttstghkn kefvarksk ldqggviqdf inalqqlsnp ellfkdwnd vraelaktpv
3601 nkkniekmye rmyaalgdpc apglgafrrk fiqtfqkefd khfgkqgskl lrmklstdfnd
3661 itnmllllkmm kdsckppgnlk ecspwmsdfk veflrnelei pgqydgrrgkp lpeyhvriag
3721 fdervtvmass lrrpkriiir ghderhpfl vkggedlrgd qrveqlfqvm ngilaqdsac
3781 sqralqlrty svvpmtssdp rappceykdw ltkmsgkhdv gaymlmykga nrtetvtser
3841 kreskvpadl lkrafvrmst speaflalrs hfasshalic ishvilgigd rhlnnfmvam
3901 etggvigidf ghafgsatqf lpvpelmpfr ltrqfinlml pmketglmys imvhalrafr
3961 sdpglltntm dvfvkepsfd wknfeqkmlk kggswiqein vaeknwyprrq kicyakrkla
4021 ganpavitcd elllghekap afirdyvavar gskdhniara epesglseet qvkcimdgat
4081 dpniltgrtwe gwepwm

2022/4

16/17

FIG. 12A

LOCUS 130781 1014 aa 01-NOV-1997
DEFINITION POLY (ADP-RIBOSE) POLYMERASE (PARP) (ADPRT)
(NAD(+)
ADP-RIBOSYLTRANSFERASE) (POLY(ADP-RIBOSE)
SYNTHETASE).
ACCESSION 130781
PID g130781
DBSOURCE SWISS-PROT: locus PPOL_HUMAN, accession P09874
class: standard.
created: Mar 1, 1989.
sequence updated: Dec 1, 1992.
annotation updated: Nov 1, 1997.
xrefs: gi: 510112, gi: 1017423, gi: 190166, gi: 190167, gi: 337423,
gi: 337424, gi: 178151, gi: 178152, gi: 190266, gi: 190267, gi:
178188, gi: 178190, gi: 189533, gi: 189534, gi: 35286, gi: 825702,
gi: 35288, gi: 189535, gi: 189536, gi: 88229, gi: 88227, gi:
627553, gi: 107162, gi: 107160, gi: 482956, gi: 420073, gi: 107158
xrefs (non-sequence databases): AARHUS/GHENT-2DPAGE 1620,
MIM
173870, MIM 173871, PROSITE PS00347, PROSITE PS50064
KEYWORDS TRANSFERASE; GLYCOSYLTRANSFERASE; NAD; DNA-
BINDING; NUCLEAR
PROTEIN; ADP-RIBOSYLATION; ZINC-FINGER; ZINC.
SOURCE human.
ORGANISM Homo sapiens
Eukaryotae; Metazoa; Chordata; Vertebrata; Mammalia; Eutheria;
Primates; Catarrhini; Hominidae; Homo.
REFERENCE 1 (residues 1 to 1014)
AUTHORS Auer,B., Nagl,U., Herzog,H., Schneider,R. and
Schweiger,M.
TITLE Human nuclear NAD+ ADP-ribosyltransferase(polymerizing):
organization of the gene
JOURNAL DNA 8 (8), 575-580 (1989)
MEDLINE 90091744
REMARK SEQUENCE FROM N.A.
REFERENCE 2 (residues 1 to 1014)
AUTHORS Uchida,K., Morita,T., Sato,T., Ogura,T., Yamashita,R.,
Noguchi,S.,
Suzuki,H., Nyunoya,H., Miwa,M. and Sugimura,T.
TITLE Nucleotide sequence of a full-length cDNA for human fibroblast
poly(ADP-ribose) polymerase
JOURNAL Biochem. Biophys. Res. Commun. 148 (2), 617-622 (1987)
MEDLINE 88076933
REMARK SEQUENCE FROM N.A.
TISSUE=FIBROBLAST

662240 " 29396660

[illegible]

1 maessdklyr veyaksgras ckkcsesipk dslrmaimvg spmfdgkvph wyhfcscfwkv
61 ghsirhpdve vdgfselrwd dqqkvkktae aggvtgkgqd gigskaektl gdfaaeyaks
121 nrstckgcme kiekqgvrls kkmvdpekpq lgmidrwyhp gcfvknreel gfrpeysasq
181 lkgfslate dkealkkqlp gvksegkrkg devdgvdeva kkkkskkekdk dsklekalka
241 qndliwnikd elkkvcstnd lkellifnkq qvpsgesail drvadgmvfg allpceecsg
301 qlvfksdayy ctgdvtawtk cmvktqtpnr kewvtpkefr eisylkkklv kkqdrifppe
361 tsasvaatpp pstasapaav nssasadkpl snmkiltlgk lsrnkdevka mieklggklt
421 gtankasici stkkevekmn kkmeevkean irvvsedflq dvsastkslq elflahilsp
481 wgaevkaepv evvaprgksg aalskkskgg vkeeginkse krmkltlkgg aavdpdsgle
541 hsahvlekkg kvfsatlglv divkgtnsyy klqlleddke nrywifrswg rvgtvigsnk
601 legmpskeda iehfmklyee ktgnawhskn ftkypkkfyp leidyggdee avkkltvnpq
661 tksklpkpqv dliknifdve smkkamveye idlqkmplgk lskrqigaay silvevqqav
721 sqgssdsqil dlsnrftytl phdfgmkkpp llnnadsvga kvemldnlll ievaysllrg
781 gsddsskdpi dvnyeklktid ikvvdrdsee aeilrkyvkn thatthnayd levidifkie
841 regecqrykp fvkqlhnrll whgsrttnfa gilsqglria ppeapvtgym fgkgiyfadm
901 vksanycht sqgdpiglil lgevalgnmy elkhashisk lpkgkhsvkg lgkttpdpsa
961 nislldqvdyv lgtgissgvn dtsllyneyi vydiaqvnlk yllklkfnfk tslw

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

DECLARATION AND POWER OF ATTORNEY

Assistant Commissioner for Patents
Washington, D.C. 20231

As a below-named inventor, I hereby declare that I believe I am:

- ☐ the original, first and sole inventor; or
- ☒ an original, first and joint inventor along with the other inventors listed below, of the subject matter which is claimed and for which a patent is sought on the invention entitled

AUTOANTIGENIC FRAGMENTS, METHODS AND ASSAYS

the specification of which ☒ is attached hereto;

☐ was filed on _____ as Application

Serial No. _____ and was amended

through _____ (if applicable).

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended as indicated above.

I acknowledge the duty to disclose to the Patent and Trademark Office all information known to me to be material to the patentability of this application in accordance with Title 37, Code of Federal Regulations, §1.56.

Foreign Priority

I hereby claim foreign priority benefits under Title 35, United States Code, §119 (a)-(d) of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate for the same invention having a filing date before that of the application on which priority is claimed:

Prior Foreign Application(s)

Priority Claimed

Country	Number	Date Filed	Attorney Docket

☐ ☐
Yes No

Country	Number	Date Filed	Attorney Docket

☐ ☐
Yes No

Prior United States Filing

I hereby claim the benefit under Title 35, United States Code, §119(e) of any United States provisional application(s) listed below.

60/082,643	April 22, 1998	20221PV
Appln. Ser. No.	Filing Date	Attorney Docket
Appln. Ser. No.	Filing Date	Attorney Docket
Appln. Ser. No.	Filing Date	Attorney Docket
Appln. Ser. No.	Filing Date	Attorney Docket

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application(s) in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, §1.56 which occurred between the filing date(s) of the prior application(s) and the national or PCT international filing date of this application:

Appln. Ser. No.	Filing Date	Status	Attorney Docket
Appln. Ser. No.	Filing Date	Status	Attorney Docket
Appln. Ser. No.	Filing Date	Status	Attorney Docket
Appln. Ser. No.	Filing Date	Status	Attorney Docket
Appln. Ser. No.	Filing Date	Status	Attorney Docket
Appln. Ser. No.	Filing Date	Status	Attorney Docket

And I hereby appoint

Michael D. Yablonsky

Jack L. Tribble

Reg. No. 40,407

Reg. No. 32,633

Reg. No. _____

respectively and individually, as my attorneys or agents with full power of substitution and revocation, to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith. Please address all communications to:

Michael D. Yablonsky
Patent Department
Merck & Co., Inc.
P.O. Box 2000
Rahway, NJ 07065-0907

Telephone No. (732) 594- 4678

I hereby declare further that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Full name of sole or joint inventor	ANTHONY ROSEN
Inventor's Signature	
Date	
Residence	Baltimore, MD 21209
Citizenship	US
Post Office Address (if different from above)	720 Rutland Avenue Baltimore, MD 21205

Full name of joint inventor	DONALD W. NICHOLSON
Inventor's Signature	
Date	
Residence	Montreal, Quebec, Canada
Citizenship	CA
Post Office Address (if different from above)	126 East Lincoln Avenue Rahway, NJ 07065-0907

Full name of joint inventor	LIVIA CASCIOLA-ROSEN
Inventor's Signature	
Date	
Residence	Baltimore, MD 21209
Citizenship	US
Post Office Address (if different from above)	720 Rutland Avenue Baltimore, MD 21205

Full name of joint inventor	FELIPE A. ANDRADE
Inventor's Signature	
Date	
Residence	Baltimore, MD 21218
Citizenship	US
Post Office Address (if different from above)	720 Rutland Avenue Baltimore, MD 21205

Full name of joint inventor	SOPHIE ROY
Inventor's Signature	
Date	
Residence	Montreal, Quebec, Canada
Citizenship	CA
Post Office Address (if different from above)	126 East Lincoln Avenue Rahway, NJ 07065-0907

Full name of joint inventor	NANCY A. THORNBERRY
Inventor's Signature	
Date	
Residence	Westfield, NJ 07090
Citizenship	US
Post Office Address (if different from above)	126 East Lincoln Avenue Rahway, NJ 07065-0907

Full name of joint inventor	
Inventor's Signature	
Date	
Residence	
Citizenship	
Post Office Address (if different from above)	

Full name of joint inventor	
Inventor's Signature	
Date	
Residence	
Citizenship	
Post Office Address (if different from above)	